arXiv:hep-th/0106094v2 9 Feb 2005

Quantum superfield supersymmetry

A.Yu. Petrov
Departamento de Físíca e Matematica,
Instituto de Físíca,
Universidade de São Paulo,
São Paulo, Brazil
and
Department of Theoretical Physics,
Tomsk State Pedagogical University
Tomsk 634041, Russia

Abstract

Superfield approach in supersymmetric quantum field theory is described. Many examples of its applications to different superfield models are considered.

1 Introduction. General properties of superspace

This paper presents itself as lecture notes in superfield supersymmetry based on lectures given at Instituto de Físíca, Universidade de São Paulo and Instituto de Físíca, Universidade Federal do Rio Grande do Sul (Porto Alegre).

The idea of supersymmetry is now considered as one of the basic concepts of theoretical high energy physics (see f.e. [1]). Supersymmetry, being a fundamental symmetry of bosons and fermions, provides possibilities to construct theories with essentially better renormalization properties since some bosonic and fermionic contributions cancel each other. Moreover, there are essentially finite supersymmetry theories **without** higher derivatives, f.e. $N = 4$ super-Yang-Mills theory. Now most specialists in quantum field theory suggest that unified theory of all interactions must be supersymmetric.

Concept of supersymmetry was introduced in known papers by Volkov and Akulov [2] and Golfand and Lichtman [3] in early 70's and received further development in [4] (the history of arising of the concept of the supersymmetry is well described in the book [5]). The essential breakthrough in supersymmetric field theory was achieved with introducing the idea of a superfield [6] (see also [7, 8]). The superfield approach in supersymmetric quantum field theory is a main topic of these lectures. We use notations introduced in [9, 10].

A superfield is a function of bosonic coordinates x^a and fermionic (Grassmann) ones $\theta^{i\alpha}, \bar{\theta}^j_{\dot{\alpha}}$. The fermionic coordinates are transformed under spinor representation of Lorentz group. The indices i, j in general case take values from 1 to N in the case of N-extended supersymmetry. Here and further we are generally interested in $N = 1$ case. However, we note that all theories with N-extended supersymmetry possess $N = 1$ formulation. The supersymmetry transformations for coordinates are

$$\delta\theta^\alpha = \epsilon^\alpha;\ \delta\bar{\theta}_{\dot{\alpha}} = \epsilon_{\dot{\alpha}};\ \delta x^a = i(\epsilon\sigma^a\bar{\theta} - \bar{\epsilon}\sigma^a\theta). \tag{1.1}$$

Here $\epsilon^\alpha, \bar{\epsilon}^{\dot{\alpha}}$ are fermionic parameters. The general form of superfield is (see f.e. [9, 11]):

$$\begin{aligned} F(x,\theta,\bar{\theta}) &= A(x) + \theta^\alpha\psi_\alpha(x) + \bar{\theta}_{\dot{\alpha}}\zeta^{\dot{\alpha}}(x) + \theta^2 F(x) + \bar{\theta}^2 G(x) + i(\bar{\theta}\sigma^a\theta)A_a(x) + \\ &+ \bar{\theta}^2\theta^\alpha\chi_\alpha(x) + \theta^2\bar{\theta}_{\dot{\alpha}}\xi^{\dot{\alpha}}(x) + \theta^2\bar{\theta}^2 H(x). \end{aligned} \tag{1.2}$$

We note that this power series is finite due to anticommutation of Grassmann numbers $\theta, \bar{\theta}$ which enforces $\theta^n, \bar{\theta}^n$ to vanish at $n \geq 3$. Further we will see that there are some restrictions on structure of superfields caused by the form of representation of supersymmetry algebra. Here $f(x), \psi_\alpha(x), \ldots$ are bosonic and fermionic fields forming component content of superfield F. If a theory describing dynamics of these fields is supersymmetric its action should be invariant under supersymmetry transformations, i.e. symmmetry transformations with fermionic parameters.

Example. In Wess-Zumino model [9] these transformations have the form

$$\begin{aligned}
\delta A(x) &= \epsilon^\alpha \psi_\alpha(x); \\
\delta \psi_\alpha(x) &= \epsilon_\alpha F(x) - \bar{\epsilon}^{\dot{\alpha}} i \partial_{\alpha\dot{\alpha}} A(x); \\
\delta F(x) &= \bar{\epsilon}_{\dot{\alpha}} i \partial^{\alpha\dot{\alpha}} \psi_\alpha.
\end{aligned} \tag{1.3}$$

Variation of arbitrary superfield $F(x, \theta, \bar{\theta})$ has the form

$$\delta F(x, \theta, \bar{\theta}) = (\epsilon^\alpha Q_\alpha + \bar{\epsilon}_{\dot{\alpha}} \bar{Q}^{\dot{\alpha}}) F(x, \theta, \bar{\theta}). \tag{1.4}$$

Here $Q_\alpha, \bar{Q}_{\dot{\alpha}}$ are generators of supersymmetry possessing anticommutation relations

$$\{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} = 2i\sigma^m_{\alpha\dot{\alpha}} \partial_m; \ \{Q_\alpha, Q_\beta\} = \{\bar{Q}_{\dot{\alpha}}, \bar{Q}_{\dot{\beta}}\} = 0; \ [Q_\alpha, \partial_m] = 0. \tag{1.5}$$

The variation (1.4) is a translation in some space.

As a result we need in introducing some extended space parametrized by bosonic and fermionic coordinates $(x^a, \theta^\alpha, \bar{\theta}^{\dot{\alpha}})$ which includes standard space-time as subspace. This extended space is called superspace. Translations on superspace are given by standard Poincare translations and transformations (1.4). It is easy to see that (1.4) is a manifestly Lorentz covariant transformation. The superspace is parametrized by 4 bosonic coordinates x^a and 4 fermionic ones $\theta^\alpha, \bar{\theta}^{\dot{\alpha}}$ so it is 8-dimensional and is denoted as $R^{4|4}$. It is natural to consider superfields as fields in the superspace. Our task is to develop quantum theory for superfields based on principles of standard quantum field theory.

To develop field theory on superspace we must introduce integration and differentiation on superspace, i.e. with respect to Grassmann coordinates. We can introduce left ∂_L and right ∂_R derivatives with respect to Grassmann coordinates as

$$\begin{aligned}
\frac{\partial_L}{\partial \theta^{\alpha_i}} (\theta^{\alpha_1} \theta^{\alpha_{i-1}} \theta^{\alpha_i} \theta^{\alpha_{i+1}} \ldots \theta^{\alpha_n}) &= (-1)^{a_1 + \ldots + a_{i-1}} (\theta^{\alpha_1} \theta^{\alpha_{i-1}} \theta^{\alpha_{i+1}} \ldots \theta^{\alpha_n}); \\
\frac{\partial_R}{\partial \theta^{\alpha_i}} (\theta^{\alpha_1} \theta^{\alpha_{i-1}} \theta^{\alpha_i} \theta^{\alpha_{i+1}} \ldots \theta^{\alpha_n}) &= (-1)^{a_{i+1} + \ldots + a_n} (\theta^{\alpha_1} \theta^{\alpha_{i-1}} \theta^{\alpha_{i+1}} \ldots \theta^{\alpha_n}).
\end{aligned} \tag{1.6}$$

Therefore these derivatives differ only by a sign factor. We can choose f.e. left one and use it henceforth.

To introduce the integral we employ the definition $\int d\theta\theta = 1$, or, generally,

$$\int d\theta^\alpha \theta_\beta = \delta^\alpha_\beta.$$

It is a convention. Note that θ and $d\theta$ have different dimensions: the mass dimension of θ is equal to $-\frac{1}{2}$, and of $d\theta$ – to $\frac{1}{2}$, and variation $\delta\theta$ cannot be mixed with differential $d\theta$. Then, integral from a constant is zero,

$$\int d\theta 1 = 0$$

this identity is caused by translation invariance due to which relation $\int d\theta(\theta+\lambda) = \int d\theta\theta$ for constant λ must be satisfied, hence $\lambda \int d\theta = 0$. We introduce the following scalar measures for Grassmann integration:

$$d^2\theta = -\frac{1}{4} d\theta^\alpha d\theta_\alpha, \; d^2\bar\theta = -\frac{1}{4} d\bar\theta_{\dot\alpha} d\bar\theta^{\dot\alpha}, \; d^4\theta = d^2\theta d^2\bar\theta. \tag{1.7}$$

These measures satisfy the relations

$$\int d^2\theta\theta^2 = \int d^2\bar\theta\bar\theta^2 = \int d^4\theta\theta^4 = 1 \tag{1.8}$$

(here and further we denote $\theta^4 \equiv \theta^2\bar\theta^2$).

Since $\frac{\partial\theta^\alpha}{\partial\theta^\beta} = \delta^\alpha_\beta$ as well as $\int d\theta^\alpha\theta_\beta = \delta^\alpha_\beta$ we conclude that integration and differentiation in Grassmann space are equivalent. F.e. we see that

$$\begin{aligned} \int d^4\theta F(x,\theta,\bar\theta) &= \frac{1}{16}\frac{\partial^2}{\partial\theta^2}\frac{\partial^2}{\partial\bar\theta^2} F(x,\theta,\bar\theta) = \frac{1}{16} F(x,\theta,\bar\theta)|_{\theta^2\bar\theta^2}; \\ \int d^2\theta G(x,\theta) &= -\frac{1}{4}\frac{\partial^2}{\partial\theta^2} G(x,\theta) = -\frac{1}{4} G(x,\theta)|_{\theta^2}. \end{aligned} \tag{1.9}$$

Here $|_{\theta^2}, |_{\theta^2\bar\theta^2}$ denotes the corresponding component of the superfield. Of course, differentiations with respect to Grassmann coordinates anticommute.

The supersymmetry generators possess several realizations in terms of $\frac{\partial}{\partial x^m}$ and $\frac{\partial}{\partial\theta_\alpha}, \frac{\partial}{\partial\bar\theta_{\dot\alpha}}$, f.e.

$$\bar Q_{\dot\alpha} = \frac{\partial}{\partial\bar\theta^{\dot\alpha}} - i\theta^\alpha(\sigma^m)_{\alpha\dot\alpha}\partial_m, \; Q_\alpha = -\frac{\partial}{\partial\theta^\alpha} + i\bar\theta^{\dot\beta}(\sigma^m)_{\dot\beta\alpha}\partial_m. \tag{1.10}$$

All possible realizations of the supersymmetry generators must satisfy relations (1.5).

The spinor supercovariant derivatives D_A also must be constructed from $\frac{\partial}{\partial x^m}$ and $\frac{\partial}{\partial\theta_\alpha}, \frac{\partial}{\partial\bar\theta_{\dot\alpha}}$. They should anticommute with generators $Q_\alpha, \bar Q_{\dot\alpha}$ which provides that $D_A\Phi$ is transformed covariantly, i.e. according to (1.4):

$$\delta(D_A\Phi) = (\epsilon Q + \bar\epsilon\bar Q) D_A\Phi.$$

F.e. if generators of supersymmetry are realized in terms of (1.10) supercovariant derivatives are realized as

$$\begin{aligned} \bar D_{\dot\alpha} &= -i\bar Q_{\dot\alpha} + i\theta^\alpha\partial_{\alpha\dot\alpha} = -i\frac{\partial}{\partial\bar\theta^{\dot\alpha}}, \\ D_\alpha &= -iQ_\alpha + i\bar\theta^{\dot\alpha}\partial_{\alpha\dot\alpha} = i(-\frac{\partial}{\partial\theta^\alpha} + 2i\bar\theta^{\dot\beta}(\sigma^m)_{\dot\beta\alpha}\partial_m). \end{aligned} \tag{1.11}$$

Here and further $\partial_{\alpha\dot{\alpha}} = (\sigma^m)_{\alpha\dot{\alpha}}\partial_m$. The spinor supercovariant derivatives satisfy the following anticommutation relations

$$\{D_\alpha, \bar{D}_{\dot{\alpha}}\} = -2i\partial_{\alpha\dot{\alpha}};\ \{D_\alpha, D_\beta\} = \{\bar{D}_{\dot{\alpha}}, \bar{D}_{\dot{\beta}}\} = 0. \tag{1.12}$$

So we defined procedures of integration and differentiation in superspace.

The next step in developing field theory is in introducing of delta function. It must satisfy the condition analogous to standard delta function

$$\int d^4\theta' \delta^4(\theta - \theta') f(\theta') = f(\theta). \tag{1.13}$$

This identity can be satisfied if we choose

$$\delta^4(\theta - \theta') = \frac{1}{16}(\theta - \theta')^2(\bar{\theta} - \bar{\theta}')^2. \tag{1.14}$$

It is easy to see that this delta function satisfies the condition

$$\int d^4\theta \delta^4(\theta - \theta') = 1. \tag{1.15}$$

We note the identity

$$\delta^4(\theta_1 - \theta_2) D_1^2 \bar{D}_2^2 \delta^4(\theta_1 - \theta_2) = 16\delta^4(\theta_1 - \theta_2). \tag{1.16}$$

Further we denote $\delta_{12} = \delta^4(\theta_1 - \theta_2)$. It is easy to see that $\delta_{12}\delta_{12} = \delta_{12}D^\alpha\delta_{12} = \delta_{12}D^2\delta_{12} = \delta_{12}\bar{D}_{\dot{\alpha}}\delta_{12} = 0$.

A supermatrix is defined as a matrix $M = M_Q^P$ of the form

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}. \tag{1.17}$$

determining a quadratic form $z_P M_Q^P z'^Q$ with z, z' are coordinates on superspace. Here A, B, C, D are even-even, even-odd, odd-even and odd-odd blocks respectively. Superdeterminant of this matrix is introduced as

$$\mathrm{sdet} M = \int d^8 z_1 d^8 z_2 exp(-z_1 M z_2). \tag{1.18}$$

It is equal to

$$\mathrm{sdet} M = \det A \det^{-1}(D - CA^{-1}B). \tag{1.19}$$

And supertrace is equal to $\mathrm{Str} M = \sum_A (-1)^{\epsilon_A} M_A^A = \mathrm{tr} A - \mathrm{tr} D$. As usual, $\mathrm{sdet} M = \exp(\mathrm{Str} \log M)$.

We can introduce change of variables in superspace. Then, if it has the form

$$x'^a = x'^a(x, \theta, \bar{\theta});\ \theta'^\alpha = \theta'^\alpha(x, \theta, \bar{\theta}),\ \bar{\theta}'^{\dot{\alpha}} = \bar{\theta}'^{\dot{\alpha}}(x, \theta, \bar{\theta}), \tag{1.20}$$

the measure of integral is transformed as

$$d^4x'd^4\theta' = d^4xd^4\theta \ \mathrm{sdet}(\frac{\partial z'}{\partial z}), \tag{1.21}$$

where supermatrix $(\frac{\partial z'}{\partial z})$ is

$$\frac{\partial z'}{\partial z} = \begin{pmatrix} \frac{\partial x'}{\partial x} & \frac{\partial x'}{\partial \theta} & \frac{\partial x'}{\partial \bar{\theta}} \\ \frac{\partial \theta'}{\partial x} & \frac{\partial \theta'}{\partial \theta} & \frac{\partial \theta'}{\partial \bar{\theta}} \\ \frac{\partial \bar{\theta}'}{\partial x} & \frac{\partial \bar{\theta}'}{\partial \theta} & \frac{\partial \bar{\theta}'}{\partial \bar{\theta}} \end{pmatrix}. \tag{1.22}$$

We also must introduce variational derivative. In common field theory it is defined as

$$\frac{\delta}{\delta A(x)} \int d^4y f(y) A(y) = f(x), \tag{1.23}$$

if $f(x)$ and $A(x)$ are functionally independent. Just analogous definition can be introduced for general (not chiral) superfield:

$$\frac{\delta}{\delta V(z)} \int d^8z' f(z') V(z') = f(z). \tag{1.24}$$

However, for chiral superfields the definition differs. Really, by definition chiral superfield $\Phi(z)$ satisfies the condition $\bar{D}_{\dot{\alpha}}\Phi = 0$. Choice of supercovariant derivatives in the form (1.11) allows one make Φ $\bar{\theta}$-independent, then the integral from a chiral function is non-trivial when it is calculated over chiral subspace, i.e. over $d^6z = d^4xd^2\theta$. Hence we must introduce variational derivative with respect to chiral superfield Φ as

$$\frac{\delta}{\delta \Phi(z)} \int d^6z' F(z') \Phi(z') = F(z). \tag{1.25}$$

And variational derivative from integral over whole superspace with respect to chiral superfield can be introduced as

$$\frac{\delta}{\delta \Phi(z)} \int d^8z' G(z') \Phi(z') = \frac{\delta}{\delta \Phi(z)} \int d^6z' (-\frac{1}{4}\bar{D}^2) G(z') \Phi(z') = -\frac{1}{4}\bar{D}^2 G(z). \tag{1.26}$$

Therefore $\frac{\delta \Phi(z)}{\delta \Phi(z')} = \delta_+(z - z')$ where $\delta_+(z - z') = -\frac{1}{4}\bar{D}^2\delta^8(z - z')$ is a chiral delta function. It allows us to obtain useful relation

$$\frac{\delta^2}{\delta\Phi(z_1)\delta\bar{\Phi}(z_2)} = \frac{1}{16}\bar{D}_1^2 D_2^2 \delta^8(z_1 - z_2) = (-\frac{1}{4})D^2\delta_+(z_1 - z_2) = (-\frac{1}{4})\bar{D}^2\delta_-(z_1 - z_2). \tag{1.27}$$

Here $\delta_-(z_1 - z_2) = -\frac{1}{4}D^2\delta^8(z_1 - z_2)$ is antichiral delta function. Note the relation $D_1^2\delta^8(z_1 - z_2) = D_2^2\delta^8(z_1 - z_2)$.

If we consider some differential operator Δ acting on superfields we can introduce its fuctional supertrace and superdeterminant:

$$\mathrm{Str}\Delta = \int d^8z_1 d^8z_2 \delta^8(z_1 - z_2) \Delta \delta^8(z_1 - z_2). \tag{1.28}$$

If we introduce kernel of the Δ which has the form $\Delta(z_1, z_2)$ we can write

$$\mathrm{Str}\Delta = \int d^8 z \Delta(z, z). \tag{1.29}$$

Superdeterminant is introduced as

$$\mathrm{sdet}\Delta = \exp \mathrm{Str}(\log \Delta). \tag{1.30}$$

Further we will be generally interested in theories describing dynamics of chiral and real scalar superfield. Note that irreducible representation of supersymmetry algebra is realized namely on these superfields [9]. The most important examples are Wess-Zumino model, general chiral superfield theory [12], $N = 1$ super-Yang-Mills theory and four-dimensional dilaton supergravity [13]. In this paper we consider application of superfield approach to these models.

2 Generating functional and Green functions for superfields

Now our aim consists of describing a method for calculation of generating functional and Green functions for superfields and following application of this method to calculation of superfield quantum corrections, i.e. in development of superfield perturbative technique. We note that during last years activity in development of nonperturbative approaches in superfield quantum theory stimulated by paper [14] essentially increased. Nevertheless perturbative approach is still the leading one, and possibility of using nonperturbative methods is frequently based on applications of the perturbative ones.

The generalization of path integral method for superfield theory turns to be quite straightforward but a bit formal. Really, generating functional is defined in terms of path integral which is well-defined only for some special cases. However, the case of Gaussian path integral is: (i) well-defined both in standard field theory and in superfield theory (ii) enough for development of superfield perturbation technique.

Let us shortly describe introduction of path integral in common field theory. Let classical action $S[\phi]$ be a local space-time functional. The equations of motion are $S_{,i}[\phi] = 0|_{\phi=\phi_0}$. The ϕ_0 is a solution for this equation. We suppose that Hessian is non-singular at this point: $\mathrm{det}S_{ij}[\phi]_{\phi=\phi_0} \neq 0$ (or as is the same equation $S_{ij}|_{\phi=\phi_0} a^j = 0$ is satisfied if and only if $a^j = 0$. If Hessian is singular, we add to the action some term to make it non-zero (in gauge theories such term is called gauge-fixing one), after adding of this term all consideration is just the same as if the Hessian is non-zero from the very beginning. We suggest that action $S[\phi]$ is analytic functional, i.e. it can be expanded into power series in a neighbourhood of ϕ_0:

$$S[\phi] = S[\phi_0] + \sum_{n=2}^{\infty} \frac{1}{n!} S_{,i_1 \ldots i_n} (\phi - \phi_0)^{i_n} \ldots (\phi - \phi_0)^{i_1}. \tag{2.1}$$

The term with $n = 2$ is called linearized action:

$$S_0 = \frac{1}{2} \tilde{\phi}^i S_{ij}[\phi_0] \tilde{\phi}^j. \tag{2.2}$$

Here and further $\tilde{\phi}^i = \phi^i - \phi^i_0$. Terms with $n \geq 3$ are called interaction terms S_{int}, and the action takes the form

$$S[\phi] = S[\phi_0] + S_0[\tilde{\phi}; \phi_0] + S_{int}[\tilde{\phi}; \phi_0]. \tag{2.3}$$

The Green function G^{ij} is determined on the base of linearized action as

$$S_{ij}[\phi_0]G^{jk} = -\delta^k_i; \; G^{ij}S_{jk}[\phi_0] = -\delta^i_k. \tag{2.4}$$

The generating functional of Green functions is introduced as

$$Z[J] = N\int D\phi \exp(\frac{i}{\hbar}(S[\phi] + J\phi)). \tag{2.5}$$

The Green functions can be obtained on the base of the generating functional as

$$< \phi(x_1)\dots\phi(x_n) >= (\frac{1}{i}\frac{\delta}{\delta J(x_1)})\dots(\frac{1}{i}\frac{\delta}{\delta J(x_n)})N\int D\phi \exp(\frac{i}{\hbar}(S[\phi] + J\phi). \tag{2.6}$$

We can calculate the path integral (2.5). To do it we expand $S[\phi] = S_0[\phi] + S_{int}[\phi]$ after changing $\tilde{\phi} \to \phi$ in (2.3), $S_0[\phi] = \int d^4x\phi\Delta\phi$ where Δ is some operator. Of course, path integration is quite formal operation well-defined only for the Gaussian integral and expressions derived from it. However, both in standard and superfield case we need mostly Gaussian integrals. As usual,

$$\begin{aligned}&\int D\phi \exp(\frac{i}{\hbar}(S[\phi] + J\phi)) = \int D\phi \exp(\frac{i}{\hbar}(\phi\Delta\phi + S_{int}[\phi] + J\phi)) = \\ = \;& \exp(\frac{i}{\hbar}S_{int}(\frac{\hbar}{i}\frac{\delta}{\delta J}))\int D\phi \exp(\frac{i}{\hbar}(\phi\Delta\phi + J\phi)).\end{aligned} \tag{2.7}$$

And (since this integral is Gaussian-like)

$$\int D\phi \exp(\frac{i}{\hbar}(\phi\Delta\phi + J\phi)) = \exp(-\frac{i}{2}J(\frac{\hbar}{\Delta})J)det^{-1/2}(\frac{\Delta}{\hbar}). \tag{2.8}$$

Therefore all dependence of sources is concentrated in $\exp(-\frac{i}{2}J(\frac{\hbar}{\Delta})J)$. Construction of Feynman diagrams from expressions (2.6, 2.7, 2.8) is quite straightforward.

Let us carry out this approach for superfield theory. Our example is Wess-Zumino model, consideration of other theories is rather analogous. We do not address specifics of gauge theories in which one must introduce gauge fixing and ghosts since after their introduction all procedure is just the same. The action of Wess-Zumino model with chiral sources is

$$S_J[\Phi, \bar{\Phi}; J, \bar{J}] = \int d^8z\Phi\bar{\Phi} + (\int d^6z(\frac{\lambda}{3!}\Phi^3 + \frac{m}{2}\Phi^2 + \Phi J) + h.c.) \tag{2.9}$$

(as usual, conjugated terms to chiral superfields are antichiral ones). It can be rewritten in terms of integrals over chiral and antichiral subspace only:

$$S_J[\Phi, \bar{\Phi}; J, \bar{J}] = \int d^6z(\frac{1}{2}\Phi(-\frac{\bar{D}^2}{4})\bar{\Phi} + \frac{\lambda}{3!}\Phi^3 + \frac{m}{2}\Phi^2 + \Phi J) + h.c. \tag{2.10}$$

The generating functional is

$$Z[J,\bar{J}] = \int D\Phi D\bar{\Phi} \exp(iS_J[\Phi,\bar{\Phi};J,\bar{J}]). \tag{2.11}$$

The action S_J (2.10) can be represented in matrix form

$$\begin{aligned} S_J &= \frac{1}{2}\int dz_1 dz_2 \begin{pmatrix} \Phi(z_1)\bar{\Phi}(z_1) \end{pmatrix} \begin{pmatrix} m & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & m \end{pmatrix} \begin{pmatrix} \delta_+(z_1-z_2) & 0 \\ 0 & \delta_-(z_1-z_2) \end{pmatrix} \times \\ &\times \begin{pmatrix} \Phi(z_2) \\ \bar{\Phi}(z_2) \end{pmatrix} + \frac{\lambda}{3!}(\int d^6 z \Phi^3 + h.c.). \end{aligned} \tag{2.12}$$

Integration in all terms is assumed with taking into account the corresponding chirality. We see that the operator Δ determining quadratic part of the action (see (2.7,2.8)) looks like

$$\Delta = \begin{pmatrix} m & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & m \end{pmatrix}. \tag{2.13}$$

The propagator is an operator inverse to this one:

$$G = \Delta^{-1} = \frac{1}{\Box - m^2}\begin{pmatrix} m & \frac{1}{4}\bar{D}^2 \\ \frac{1}{4}D^2 & m \end{pmatrix}. \tag{2.14}$$

In other words, propagator G satisfies the equation

$$\Delta G = -\begin{pmatrix} \delta_+(z_1-z_2) & 0 \\ 0 & \delta_-(z_1-z_2) \end{pmatrix}. \tag{2.15}$$

The matrix $\mathbf{1} = \begin{pmatrix} \delta_+(z_1-z_2) & 0 \\ 0 & \delta_-(z_1-z_2) \end{pmatrix}$ plays the role of functional unit matrix.

Thus, the generating functional is

$$\begin{aligned} Z[J,\bar{J}] &= \exp(i\frac{\lambda}{3!}\int d^6 z(\frac{\delta}{\delta J(z)})^3 + h.c.) \overset{-1/2}{\det} \Delta \times \\ &\times \exp\Big\{ -\frac{i}{2}\int dz_1 dz_2 \begin{pmatrix} J(z_1)\bar{J}(z_1) \end{pmatrix} \frac{1}{\Box - m^2} \begin{pmatrix} m & \frac{1}{4}\bar{D}^2 \\ \frac{1}{4}D^2 & m \end{pmatrix} \times \\ &\times \begin{pmatrix} \delta_+(z_1-z_2) & 0 \\ 0 & \delta_-(z_1-z_2) \end{pmatrix} \begin{pmatrix} J(z_2) \\ \bar{J}(z_2) \end{pmatrix} \Big\}. \end{aligned} \tag{2.16}$$

The argument of the exponential function in last expression can be rewritten as

$$-\frac{i}{2}\Big(\int d^6 z J \frac{m}{\Box - m^2} J + 2\int d^6 z J \frac{\frac{1}{4}\bar{D}^2}{\Box - m^2}\bar{J} + \int d^6 \bar{z} \bar{J} \frac{m}{\Box - m^2}\bar{J}\Big). \tag{2.17}$$

We can introduce two-point Green functions:

$$\begin{aligned} G_{++}(z_1,z_2) &= \frac{1}{i^2}\frac{\delta^2 Z[J]}{\delta J(z_1)\delta J(z_2)} = (-\frac{1}{4})^2 \bar{D}_1^2 \bar{D}_2^2 K_{++}(z_1,z_2) \\ G_{+-}(z_1,z_2) &= \frac{1}{i^2}\frac{\delta^2 Z[J]}{\delta J(z_1)\delta \bar{J}(z_2)} = (-\frac{1}{4})^2 \bar{D}_1^2 D_2^2 K_{+-}(z_1,z_2) \\ G_{--}(z_1,z_2) &= \frac{1}{i^2}\frac{\delta^2 Z[J]}{\delta \bar{J}(z_1)\delta \bar{J}(z_2)} = (-\frac{1}{4})^2 D_1^2 D_2^2 K_{--}(z_1,z_2). \end{aligned} \tag{2.18}$$

Here $K_{+-}(z_1,z_2)=K_{-+}(z_1,z_2)=-\frac{1}{\Box-m^2}\delta^8(z_1-z_2)$, $K_{++}(z_1,z_2)=\frac{m\bar{D}^2}{4\Box(\Box-m^2)}\delta^8(z_1-z_2)$, $K_{--}(z_1,z_2)=\frac{mD^2}{4\Box(\Box-m^2)}\delta^8(z_1-z_2)$. We note that in theory of standard (not chiral) superfield the variational derivatives with respect to sources do not involve factors D^2, $\bar{D}^2$. These factors are caused by chirality. F.e. for theory of real scalar superfield with the action $\frac{1}{2}\int d^8zv\Box v$ the propagator is simply $G(z_1,z_2)=\frac{1}{\Box}\delta(z_1-z_2)$.

Different vacuum expectations can be expressed in terms of the generating functional (2.16) as

$$\begin{aligned}
&< \phi(x_1)\ldots\phi(x_n)\bar{\phi}(y_1)\ldots\bar{\phi}(y_m) >= \\
= \ & (\frac{1}{i}\frac{\delta}{\delta J(x_1)})\ldots(\frac{1}{i}\frac{\delta}{\delta J(x_n)})(\frac{1}{i}\frac{\delta}{\delta \bar{J}(y_1)})\ldots(\frac{1}{i}\frac{\delta}{\delta \bar{J}(y_m)})\times \\
\times \ & \exp\Big\{i\frac{\lambda}{3!}\int d^6z(\frac{\delta}{\delta J(z)})^3+h.c.)\overset{-1/2}{\det}\Delta\times \\
\times \ & \exp(-\frac{i}{2}\int dz_1dz_2\begin{pmatrix} J(z_1)\bar{J}(z_1)\end{pmatrix}\frac{1}{\Box-m^2}\begin{pmatrix} m & \frac{1}{4}\bar{D}^2 \\ \frac{1}{4}D^2 & m\end{pmatrix}\times \\
\times \ & \begin{pmatrix}\delta_+(z_1-z_2) & 0 \\ 0 & \delta_-(z_1-z_2)\end{pmatrix}\begin{pmatrix} J(z_2) \\ \bar{J}(z_2)\end{pmatrix}\Big\}.
\end{aligned}\tag{2.19}$$

Of course, this expression contains all orders in coupling λ. To obtain vacuum expectations up to some order in couplings we should expand $\exp(i\frac{\lambda}{3!}\int d^6z(\frac{\delta}{\delta J(z)})^3+h.c.)$ into power series. As a result as usual we arrive at some Feynman diagrams. In these diagrams $n+m$ is the number of external points, and order in λ is the number of internal points. Each vertex evidently corresponds to integration over d^6z or $d^6\bar{z}$. Therefore we have to introduce diagrams for superfield theory, i.e. Feynman supergraphs. Their value consists of the fact that they allow one to preserve manifest supersymmetry covariance at any step of calculations.

Generating functionals of arbitrary models can be constructed by analogy with Wess-Zumino model:

$$Z[\vec{J}]=\exp(i(S[\vec{\phi}]+\vec{\phi}\vec{J})).\tag{2.20}$$

Here $\vec{\phi}$ is a column matrix denoting set of all superfields, $\vec{J}$ is a column matrix denoting set of corresponding sources. The Green functions can be determined in analogy with (2.19).

3 Feynman supergraphs

It is easy to see that the Green functions (2.18), the generating functional (2.16) and the vacuum expectations (2.19) lead to the known supergraph technique. Really, any $<\phi\bar{\phi}>$-propagator corresponds to $(\Box-m^2)^{-1}$, at a chiral vertex each propagator is associated with $-\frac{1}{4}\bar{D}^2$, and at an antichiral one – with $-\frac{1}{4}D^2$. However, each chiral (antichiral) vertex corresponds to integration over d^6z ($d^6\bar{z}$). However, since we deal with $\delta^8(z_1-z_2)$, for sake of unity it is more convenient to represent all contributions in the form of integrals over d^8z via the rule $d^6z(-\frac{1}{4})\bar{D}^2F = \int d^8zF$. As a result, $\int d^6z\Phi^n$-vertex is associated with $n-1$ $(-\frac{1}{4}\bar{D}^2)$ factors, and $\int d^6\bar{z}\bar{\Phi}^m$-vertex – with $m-1$ $(-\frac{1}{4}D^2)$-factors – of course, in the case when all superfields are contracted into propagators. And the vertex $d^8z\Phi^m\bar{\Phi}^n$ in the same case – with m factors and n $(-\frac{1}{4}D^2)$ factors. Here and further we refer to superfields contracted into propagators as to the quantum ones. We see that the number of D^2, $\bar{D}^2$ factors for such vertices is number of antichiral (chiral) **quantum** superfields associated with this vertex. There is no $D, \bar{D}$-factors arisen from propagators of non-chiral (f.e. real) superfields. The propagator $<\phi\phi>$ $(<\bar{\phi}\bar{\phi}>)$ corresponds to $\frac{m\bar{D}^2}{4\Box(\Box-m^2)}$ $(\frac{mD^2}{4\Box(\Box-m^2)})$. However, only quantum fields (i.e. those ones contracted into propagators) correspond to D^2, $\bar{D}^2$ factors. External lines do not carry such a factor, and if one, two... n chiral (antichiral) superfields associated with the vertex are external the number of $\bar{D}^2$ (D^2) factors corresponding to this vertex is less by one, two... n than in the case when all superfields are contracted to propagators.

If we consider the theory of $N=1$ super-Yang-Mills (SYM) field, its quadratic action after gauge fixing looks like

$$S = \frac{1}{2}tr\int d^8zV\Box V. \tag{3.1}$$

The propagator is

$$G(z_1,z_2) = \frac{1}{\Box}\delta^8(z_1-z_2) \tag{3.2}$$

(note the opposite sign with respect to $<\phi\bar{\phi}>$-propagator). Here tr is matrix trace (the superfield V is Lie-algebra valued). There is no D factors associated with this propagator but they are associated with vertices. In pure $N=1$ SYM theory vertices are given by

$$S_{int} = \frac{g}{8}\int d^8z(\bar{D}^2D^\alpha V)[V, D_\alpha V] + \dots \tag{3.3}$$

Here dots denote higher orders in V. The vertices of any order involve are two D factors and two $\bar{D}$ factors. The D-factors in vertices involving both real and chiral (antichiral) superfields are arranged in a common way, i.e. any vertex $\Phi\bar{\Phi}V^n$ involves one factor $(-\frac{1}{4}\bar{D}^2)$ acting to the Φ superfield when it is contracted to $<\Phi\bar{\Phi}>$ propagator and one factor $(-\frac{1}{4}D^2)$ acting to the $\bar{\Phi}$ superfield when it is contracted to $<\Phi\bar{\Phi}>$ propagator. As a result we can formulate Feynman rules.
Propagators look like

$$\begin{aligned} <\phi\bar{\phi}> &= -\frac{1}{\Box-m^2}\delta^8(z_1-z_2); \\ <\phi\phi> &= \frac{m\bar{D}^2}{\Box(\Box-m^2)}\delta^8(z_1-z_2); \\ <vv> &= \frac{1}{\Box}\delta^8(z_1-z_2); \end{aligned} \tag{3.4}$$

vertices (here ϕ, $\bar{\phi}$ are quantum superfields) correspond to

$$\begin{aligned} \int d^6 z \phi^n &\rightarrow (n-1)(-\frac{1}{4})\bar{D}^2; \\ \int d^8 z \phi\bar{\phi} v^m &\rightarrow (-\frac{1}{4})D^2(-\frac{1}{4})\bar{D}^2. \end{aligned} \tag{3.5}$$

All derivatives in derivative depending vertices act on the propagators. Any external chiral (antichiral) fields do not correspond to $\bar{D}\,(D)$-factors.

Of course, it is more suitable to make Fourier representation for all propagators (note that Fourier transformation is carried out with respect to bosonic coordinates only) by the rule

$$\tilde{f}(k) = \int \frac{d^4k}{(2\pi)^4} f(x) e^{ikx}. \tag{3.6}$$

The propagators in momentum representation look like

$$< \phi(1)\bar{\phi}(2) > \;=\; \frac{1}{k^2+m^2}\delta^4_{12}; \tag{3.7}$$

$$\begin{aligned} < \phi(1)\phi(2) > &= \frac{m\bar{D}^2}{4k^2(k^2+m^2)}\delta^4_{12}; \\ < v(1)v(2) > &= -\frac{1}{k^2}\delta^4_{12}. \end{aligned} \tag{3.8}$$

Here 1, 2 are numbers of arguments, and $\delta^4_{12} \equiv \delta^4(\theta_1-\theta_2) = \frac{1}{16}(\theta_1-\theta_2)^2(\bar{\theta}_1-\bar{\theta}_2)^2$ is a Grassmann delta function. The D-factors are introduced as above. Note, however, that spinor derivatives depend after Fourier transform on momentum of propagator with which they are associated. The external superfields also can be represented in the form of Fourier integral. Each propagator is parametrized by momentum, and any vertex corresponds to integration over $d^4\theta$, coupling and delta function over incoming momenta. As usual, contribution of supergraph includes integration over all momenta and combinatoric factor which is totally analogous to that one in standard quantum field theory.

Essentially new feature of superfield theories is presence of D-factors. To evaluate D-algebra we can transport them via integration by parts, then, we can use the identity

$$\delta^4_{12} D^2 \bar{D}^2 \delta^4_{12} = 16\delta^4_{12} \tag{3.9}$$

To prove this identity we can use expansion of supercovariant derivatives (1.11) and note that due to the evident property
$\delta^4_{12}\frac{\partial}{\partial\theta^\alpha}\delta^4_{12} = \frac{\partial}{\partial\theta^\alpha}\delta^4_{12}|_{\theta_1=\theta_2} = \frac{1}{8}(\theta_{1\alpha}-\theta_{2\alpha})(\bar{\theta}_1-\bar{\theta}_2)^2|_{\theta_1=\theta_2} = 0$
only terms of the form $\delta^4_{12}(\frac{\partial}{\partial\theta})^2(\frac{\partial}{\partial\bar{\theta}})^2\delta^4_{12} = 16\delta^4_{12}$ survive.

We can prove the following **theorem**.

The final result for the contribution of any supergraph should have the form of **one** integral over $d^4\theta$.

Proof: Let us consider propagator with L loops, V vertices and P propagators. Any vertex contains integration over $d^4\theta$, i.e. there are V such integrations. Then, due to

(3.7) any propagator carries a delta function over Grassmann coordinates, i.e. there are P delta functions. Then, in any loop we can reduce the number of delta functions by one using identity (3.9), i.e. there are $P-L$ independent delta functions. As a result we can carry out $P-L$ integrations from V, and after D-algebra transformations we stay with $V-(P-L)$ integrations. And $V-(P-L)=1$, therefore the result contains one integration over $d^4\theta$. The theorem is proved [15].

This theorem is often called non-renormalization theorem. It means that all quantum corrections are local in θ-space. This theorem is often naively treated as a proof of absence of chiral corrections (proportional to integral over $d^2\theta$). However, such interpretation is wrong since any contribution in the form of integral over chiral subspace can be rewritten as an integral over whole superspace using identity

$$\int d^6z f(\Phi) = \int d^8z(-\frac{D^2}{4\Box})f(\Phi) \tag{3.10}$$

(this observation was firstly made in [16], its consequences will be studied further).

Now let us study evaluation of contributions from supergraphs. The algorithm of it is the following one.

1. We start with one of loops. If the number of D-factors in this loop is equal to 4 we turn to step 2. If it is more than 4, superfluous D-factors can be transported to external lines or another loops via integration by parts, and some of them are converted into internal momenta via identities $D^2\bar{D}^2D^2=16\Box D^2, \{D_\alpha,\bar{D}_\beta\}=-2i\partial_{\alpha\beta}$. As a result we stay with exactly 4 D-factors. If the number of D-factors is less than 4 than contribution from the entire supergraph is equal to zero.

2. We contract this loop into a point using identity (3.9) and intergrate over one of $d^4\theta$ via delta function free of derivatives.

3. This procedure is repeated for next loops.

4. We integrate over internal momenta.

However, the best way to study evaluation of supergraphs is in considering some examples.

Example 1. One-loop supergraph in Wess-Zumino model.



Fig.1

The contribution of this supergraph is equal to

$$\begin{aligned} I_1 &= \frac{1}{2}\lambda^2\int d^4\theta_1 d^4\theta_2\int\frac{d^4p}{(2\pi)^4}\Phi(-p,\theta_1)\bar{\Phi}(p,\theta_2)\delta^4_{12}\frac{\bar{D}^2_1D^2_2}{16}\delta^4_{12}\times \\ &\times \int\frac{d^4k}{(2\pi)^4}\frac{1}{(k^2+m^2)((k+p)^2+m^2)} \end{aligned} \tag{3.11}$$

The number of D-factors is just 4. D-algebra transformations are trivial: we use identity (3.9) and write $\delta^4_{12}\frac{\bar{D}_1^2 D_2^2}{16}\delta^4_{12} = \delta^4_{12}$. The free delta function δ^4_{12} allows us to integrate over $d^4\theta_2$ and denote $\theta_1 = \theta$. As a result we get

$$I_1 = \frac{1}{2}\lambda^2 \int d^4\theta \int \frac{d^4p}{(2\pi)^4}\Phi(-p,\theta)\bar{\Phi}(p,\theta)\int \frac{d^4k}{(2\pi)^4}\frac{1}{(k^2+m^2)((k+p)^2+m^2)} \qquad (3.12)$$

Integral over k can be calculated via dimensional regularization, the result for it is

$$\int \frac{d^4k}{(2\pi)^4}\frac{1}{(k^2+m^2)((k+p)^2+m^2)} = \frac{1}{16\pi^2}(\frac{1}{\epsilon} - \int_0^1 dt \log\frac{p^2t(1-t)+m^2}{\mu^2}) \qquad (3.13)$$

As a result, contribution of this supergraph takes the form

$$I_1 = \frac{1}{2}\lambda^2 \int d^4\theta \int \frac{d^4p}{(2\pi)^4}\Phi(-p,\theta)\bar{\Phi}(p,\theta)\frac{1}{16\pi^2}(\frac{1}{\epsilon} - \int_0^1 dt \log\frac{p^2t(1-t)+m^2}{\mu^2}) \qquad (3.14)$$

However, the regularization in superfield theory in higher loops possesses some peculiarities [17].

Example 2. Two-loop supergraph in Wess-Zumino model.



Fig.2

The contribution of this supergraph is equal to

$$\begin{aligned} I_2 &= \frac{\lambda^2}{6}\int \frac{d^4k d^4l}{(2\pi)^8}\int d^4\theta_1 d^4\theta_2 (-\frac{D_1^2}{4})\delta^4_{12}\frac{D_1^2\bar{D}_2^2}{16}\delta^4_{12}(-\frac{\bar{D}_2^2}{4})\delta^4_{12} \times \\ &\times \frac{1}{(k^2+m^2)(l^2+m^2)((k+l)^2+m^2)} \end{aligned} \qquad (3.15)$$

First we do D-algebra transformations: we can write

$$(-\frac{D_1^2}{4})\delta^4_{12}\frac{D_1^2\bar{D}_2^2}{16}\delta^4_{12}(-\frac{\bar{D}_2^2}{4})\delta^4_{12} = \delta^4_{12}\frac{D_1^2\bar{D}_2^2}{16}\delta^4_{12}\frac{D_1^2\bar{D}_2^2}{16}\delta^4_{12}$$

Then we use identity (3.9) two times:

$$\delta^4_{12}\frac{D_1^2\bar{D}_2^2}{16}\delta^4_{12}\frac{D_1^2\bar{D}_2^2}{16}\delta^4_{12} = \delta^4_{12}$$

As a result we can integrate over θ_2 using the delta function. We get

$$I_2 = \frac{\lambda^2}{6}\int \frac{d^4k d^4l}{(2\pi)^8}\int d^4\theta_1 \frac{1}{(k^2+m^2)(l^2+m^2)((k+l)^2+m^2)} \qquad (3.16)$$

This integral vanishes in the standard case since it is proportional to an integral over $d^4\theta$ from constant. However, if we suppose that m is not a constant but θ-dependent superfield this contribution is not zero. Namely this case is studied when the effective action is studied and m is suggested to depend on background superfields.

Example 3. One-loop supergraph in dilaton supergravity.


$G(k)$
$\bar{D}^{\dot{\beta}}D^2$
$\bar{D}^2D^{\alpha}$
$\partial_{\beta\dot{\beta}}$
$\partial_{\alpha\dot{\alpha}}$
$\bar{D}^2D^{\beta}$
$D^2\bar{D}^{\dot{\alpha}}$
$G(k+p)$

Fig.3

The contribution of this supergraph is equal to

$$\begin{aligned} I_3 &= \frac{\xi_1^2}{2}\int d^4\theta_1 d^4\theta_2 \int \frac{d^4p}{(2\pi)^4}\frac{d^4k}{(2\pi)^4}(\partial_{\alpha\dot{\alpha}}\sigma(-p,\theta_1))(\partial_{\beta\dot{\beta}}\sigma(p,\theta_2)) \times \\ &\times \frac{D^{\alpha}\bar{D}^2D^2\bar{D}^{\dot{\beta}}}{16}\delta^4_{12}\frac{\bar{D}^{\dot{\alpha}}D^2\bar{D}^2D^{\beta}}{16}\delta^4_{12}G(k)G(k+p). \end{aligned} \qquad (3.17)$$

Here $G(k), G(k+p)$ are functions of momenta which explicit form is not essential here (they are exactly found in [13]). The derivatives $\partial_{\alpha\dot{\alpha}}, \partial_{\beta\dot{\beta}}$ are not transported from external fields $\sigma, \bar{\sigma}$. Our aim here is to obtain terms proportional to $\partial^m\sigma\partial^n\bar{\sigma}$. We suggest that spinor derivatives associated with one propagator depend on momentum k, and with another – to $k+p$.

Using commutation relations (1.12) we find that

$$\frac{D^{\alpha}\bar{D}^2D^2\bar{D}^{\dot{\beta}}}{16}\delta^4_{12}\frac{\bar{D}^{\dot{\alpha}}D^2\bar{D}^2D^{\beta}}{16}\delta^4_{12} = \frac{2k^{\alpha\dot{\gamma}}\bar{D}_{\dot{\gamma}}D^2\bar{D}^{\dot{\beta}}}{16}\delta^4_{12}\frac{2(k+l)^{\gamma\dot{\alpha}}D_{\gamma}\bar{D}^2D^{\beta}}{16}\delta^4_{12}. \qquad (3.18)$$

We transport all spinor supercovariant derivatives to one propagator (here the terms with spinor supercovariant derivatives moved to the external lines are omitted as the irrelevant ones since they do not contribute to the divergent part [13]). As a result we arrive at

$$4k^{\alpha\dot{\gamma}}(k+p)^{\gamma\dot{\alpha}}\delta^4_{12}\frac{\bar{D}^{\dot{\beta}}D^2\bar{D}_{\dot{\gamma}}D_{\gamma}\bar{D}^2D^{\beta}}{256}\delta^4_{12}. \qquad (3.19)$$

We can use (1.12) several times. At the end we get

$$4k^{\alpha\dot{\gamma}}(k+p)^{\gamma\dot{\alpha}}(k+p)_{\gamma\dot{\gamma}}(k+p)^{\dot{\beta}\delta}\delta^4_{12}D_{\delta}\bar{D}^2D^{\beta}\delta^4_{12}. \qquad (3.20)$$

Equations (1.12) and (3.9) allow one to write

$$\delta^4_{12}D_{\delta}\bar{D}^2D^{\beta}\delta^4_{12} = -\frac{1}{2}16\delta^{\beta}_{\delta}\delta^4_{12}.$$

We substitute this expression in (3.18). Using identity $k^{\alpha\dot{\beta}}k_{\gamma\dot{\beta}} = \delta^{\alpha}_{\gamma}k^2$ we obtain the contribution from (3.18) in the form

$$64k^{\alpha\dot{\alpha}}(k+p)^{\beta\dot{\beta}}(k+p)^2\delta^4_{12},$$

which after integration over θ_2 leads to the following contribution for I_3:

$$\begin{aligned} I_3 &= 64\frac{\xi_1^2}{2}\int d^4\theta_1\int\frac{d^4p}{(2\pi)^4}\frac{d^4k}{(2\pi)^4}(\partial_{\alpha\dot\alpha}\sigma)(-p,\theta_1)(\partial_{\beta\dot\beta}\sigma)(p,\theta_1)\times \\ &\times\; k^{\alpha\dot\alpha}(k+p)^{\beta\dot\beta}(k+p)^2G(k)G(k+p). \end{aligned} \tag{3.21}$$

Detailed analysis carried out in [13] shows that this correction is divergent.

Calculation of corrections from supergraphs in other superfield theories is carried out on the base of analogous approach.

We demonstrated that supergraph technique is a very efficient method for consideration of quantum corrections in superfield theories whereas the component study is much more complicated since one supergraph corresponds to several component diagrams (it is amusing that the exact expression for the classical action of dilaton supergravity occupies a whole page [13]). The next step of its development is introducing renormalization in these theories.

4 Superficial degree of divergence. Renormalization.

We found that divergent quantum corrections arise in superfield theories as well as in standard field theories. Therefore we face two problems:

(i) to classify possible divergences;

(ii) to develop a procedure of renormalization in superfield theories.

It turns out that the technique for solving these problems is quite analogous to that one used in standard field theory. First problem can be solved on the base of superficial degree of divergence. The natural way for solving second one is in introducing superfield counterterms which are quite analogous to standard ones.

First of all let us consider superficial degree of divergence [18].

Example. The $N=1$ super-Yang-Mills (SYM) theory with chiral matter (with Wess-Zumino self-interaction). For all other models consideration is quite analogous. The action of the theory is

$$\begin{aligned} S &= \int d^8z\bar\Phi_i(e^{gV})^i_j\Phi^j+(\int d^6z(\frac{1}{2}m_{ij}\Phi_i\Phi_j+\frac{\lambda_{ijk}}{3!}\Phi_i\Phi_j\Phi_k)+h.c.)+ \\ &+\; \mathrm{tr}\frac{1}{g^2}\int d^8z(e^{-gV}D^\alpha e^{gV})\bar D^2(e^{-gV}D_\alpha e^{gV}). \end{aligned} \tag{4.1}$$

The lower-order vertices in this theory are

$$\frac{\lambda_{ijk}}{3!}\int d^6z\Phi_i\Phi_j\Phi_k,\ \frac{\lambda_{ijk}}{3!}\int d^6\bar z\bar\Phi_i\bar\Phi_j\bar\Phi_k,\ \int d^8z\bar\Phi_iV^i_j\Phi^j,\ \frac{1}{2}\mathrm{tr}\int d^8z(\bar D^2D^\alpha V)[V,D_\alpha V] \tag{4.2}$$

and higher ones. Indices i,j are matrix indices since Φ_i is an isospinor, and $V\equiv V^A(T^A)^i_j$ is Lie-algebra valued. However, all vertices corresponding to pure SYM self-interaction contain exactly two chiral and two antichiral derivatives. We have proved already that all corrections should be proportional to one integral over $d^4\theta$.

As usual, the superficial degree of divergence (SDD) is the order of the integral over internal momenta for corresponding contribution, or, as is the same, as a degree of homogeneity of diagram in momenta, considered after evaluation of D-algebra transformations [10]. The only difference of the SDD in our case is the additional impact from D-factors.

It is easy to see that contributions to the SDD are generated by momentum depending factors in propagators and vertices (as usual, any internal momentum k gives contribution 1), loop integrations, or, in other words, by manifest momentum dependence which is associated with propagators and loop integration, and by D-factors which are associated with propagators and vertices (note that due to identities $D^2\bar{D}^2D^2 = 16\Box D^2$, $\{D_\alpha, \bar{D}_{\dot{\alpha}}\} = -2i\partial_{\alpha\dot{\alpha}}$ one chiral derivative combined with an antichiral one can be converted to one momentum; therefore any D-factor contribute to the SDD with 1/2). If not all spinor derivatives are converted to internal momenta, the SDD from supergraph evidently decreases.

Let us consider arbitrary supergraph with L loops, V vertices, P propagators (C of them are $<\phi\phi>$, $<\bar{\phi}\bar{\phi}>$ -propagators) and E external lines (E_c of them are chiral). We denote the SDD as ω.

Any integration over internal momentum (i.e. over d^4k) contributes to SDD with 4. Since the number of integrations over internal momenta is the number of loops, the total contribution from all such integrations is $4L$. Any propagator includes $\frac{1}{k^2+m^2}$ or $\frac{1}{k^2}$ (3.7), hence contribution of all propagators is equal to $-2P$. Since $<\Phi\Phi>$, $<\bar{\Phi}\bar{\Phi}>$-propagator contains additional $\frac{1}{k^2}$ these propagators give additional contribution $-2C$. Therefore manifest dependence of momenta gives contribution to ω equal to $4L - 2P - 2C$.

Now let us consider contribution of D-factors to SDD. Each **vertex** (both pure gauge one and that one containing chiral superfields) without external chiral (antichiral) lines contains four D-factors (4.2) since any superfield ϕ (contracted to propagator) corresponds to $\bar{D}^2$, and $\bar{\phi}$ – to D^2. Therefore each vertex gives contribution 2. However, external chiral (antichiral) lines do not correspond to D-factors. As a result, any external line decreases ω by 1, Each $<\phi\phi>$, $<\bar{\phi}\bar{\phi}>$-propagator contains a factor $\bar{D}^2$ (D^2) with contribution 1. Then, due to identity (3.9) contraction of any loop into a point decreases the number of D-factors which can be converted to internal momenta by 4, and ω – by 2. As a result the total contribution of D-factors to ω is equal to $2V - E_c - 2L + C$ (remind that each D-factor contributes to ω with 1/2).

Therefore SDD is equal to

$$\omega = 4L - 2P - 2C + 2V - E_c - 2L + C = 2L - 2P + 2V - C - E_c. \tag{4.3}$$

Using known topological identity $L + V - P = 1$ we have

$$\omega = 2 - C - E_c. \tag{4.4}$$

Really, the SDD can be lower than (4.4) if some of D-factors are transported to external lines and do not generate internal momenta. If N_D D-factors are moved to external lines the ω is equal to

$$\omega = 2 - C - E_c - \frac{1}{2}N_D. \tag{4.5}$$

This is the final expression for the SDD. As usual, at $\omega \geq 0$ supergraph diverges, and at $\omega < 0$ – converges. We note that:
1. $\omega \leq 2$ hence SDD is restricted from above.
2. As the number of external lines grows, ω decreases. Therefore the number of divergent structures is essentially restricted – it is finite (really, there can be no more than two external chiral legs and no more than two $<\Phi\Phi>, <\bar{\Phi}\bar{\Phi}>$ propagators). And if the number of divergent structures is finite the theory is renormalizable. Hence we shown that the theory including chiral superfields with Wess-Zumino-type interaction and gauge superfields with action (4.1) is renormalizable. This is quite natural since the mass dimension of all couplings in this theory is zero.

However, non-renormalizable superfield theories also exist.

Example. General chiral superfield model [12].

The action of the model is

$$\begin{aligned} S &= \int d^8 z K(\Phi,\bar{\Phi}) + (\int d^6 z W(\Phi) + h.c.) = \\ &= \int d^8 z \Phi\bar{\Phi} + [\int d^6 z (\frac{1}{2} m\Phi^2 + \frac{\lambda}{3!}\Phi^3) + h.c.] + \\ &+ \int d^8 z [K_{12}\Phi\bar{\Phi}^2 + K_{21}\bar{\Phi}\Phi^2 + \sum_{m,n=2}^{\infty} \frac{1}{m!n!} K_{mn}\Phi^n\bar{\Phi}^m] + (\int d^6 z \sum_{l=4}^{\infty} \frac{W_n}{n!}\Phi^n + h.c.). \end{aligned} \tag{4.6}$$

Here K_{ij}, W_l are constants.

Propagators in the theory are just (3.7), their contribution to SDD is equal to $4L-2P-2C$ as above. However, the contribution from D-factors differs. Any vertex $K_{nm}\Phi^n\bar{\Phi}^m$ corresponds to n $\bar{D}^2$-factors and m D^2-factors. The total contribution to ω from all such vertices is $\sum_{V_t}(n_v+m_v)$, i.e. sum of n and m over all vertices corresponding to integral over total superspace. Any vertex $W_l\Phi^l$ contains an integral over $d^6 z$ and effectively corresponds to $(l-1)$ $\bar{D}^2$-factors. Total contribution from such vertices is $\sum_{V_c}(l_c-1)$ (i.e. sum over all purely chiral or antichiral vertices). Again external lines decrease the number of D^2 ($\bar{D}^2$)-factors by $2E_c$ (E_c is a number of external lines), each $<\Phi\Phi>, <\bar{\Phi}\bar{\Phi}>$-propagators carries one $\bar{D}^2$ (D^2)-factor. Contraction of each loop to a point decreases the number of D-factors by 4. Hence the total number of D-factors is

$$2\sum_{V_t}(n_v+m_v) + 2\sum_{V_c}(l_c-1) - 2E_c - 4L + 2C. \tag{4.7}$$

Contribution to SDD from D-factors is their number divided by two. Therefore total SDD is equal to

$$\begin{aligned} \omega &= 4L - 2P - 2C + \frac{1}{2}(2\sum_{V_t}(n_v+m_v) + 2\sum_{V_c}(l_c-1) - 2E_c - 4L + 2C) = \\ &= 2 - 2V - C - 2E_c + [\sum_{V_t}(n_v+m_v) + \sum_{V_c}(l_c-1)]. \end{aligned} \tag{4.8}$$

Here we used $2L-2P = 2-2V$. However, any vertex gives contribution -2 to term $-2V$ and l_c-1 or n_v+m_v to other terms of ω. It is evidently that either l_c-1 or

$n_v + m_v$ can be more than 2 since either $l_c \geq 3$ or $n_v + m_v \geq 3$. Hence in general case $\sum_{V_t}(n_v+m_v)+\sum_{V_c}(l_c-1)-2V \geq 0$, the number of divergent structures is not restricted, and the theory is non-renormalizable. This is quite natural since constants K_{ij} (if i or j no less than 2) and W_l (if $l \geq 4$) have negative mass dimension.

The next problem is introduction of regularization. The most natural way of introducing regualarization in supersymmetric theories is dimensional regularization. It can be introduced as usual: integral

$$\int \frac{d^4k}{(2\pi)^4}\frac{1}{(k^2+m^2)^N}$$

is replaced by

$$\int \frac{d^{4+\epsilon}k}{(2\pi)^{4+\epsilon}}\frac{1}{(k^2+m^2)^N}.$$

All divergences corresponds to poles in ϵ (no more than $\frac{1}{\epsilon^L}$ for L-loop correction).

However, there are some peculiarities. First of all, at component level any supersymmetric action includes spinors and hence γ-matrices which are well defined if and only if the dimension of space-time is integer. Therefore we must use some modification of the dimensional regularization called dimensional reduction. According to it all objects with well behaviour only at separate dimensions (such as spinors and γ-matrices) are evaluated at these dimensions (or namely at dimension equal to 4), and integrals over momenta – at arbitrary dimension. However, dimensional reduction leads to some difficulties in calculation of higher loop corrections since many supergraphs involve contractions of essentially four-dimensional objects, such as Levi-Civita tensor ϵ^{abcd}, with d-dimensional objects, and such contractions need additional definition. As a result frequently the ambiguities arise. However, such phenomena are observed only beyond two loops.

We also can use analytic regularization which corresponds to change

$$\int \frac{d^4k}{(2\pi)^4}\frac{1}{(k^2+m^2)^n} \to \int \frac{d^4k}{(2\pi)^4}\frac{1}{(k^2+m^2)^{n+\epsilon}}.$$

However, this regularization also leads to some difficulties (see discussion of questions connected to regularization in supersymmetric theories in [17]).

Technique for renormalization in superfield theories is quite analogous to that one in common QFT. It is carried out via introduction of counterterms.

Example. Consider one-loop contribution to the kinetic term in Wess-Zumino model. Corresponding supergraph is given by Fig. 1 (see above), its contribution is equal to

$$I_1 = \frac{1}{2}\lambda^2\int d^4\theta\int\frac{d^4p}{(2\pi)^4}\Phi(-p,\theta)\bar{\Phi}(p,\theta)\frac{1}{16\pi^2}(\frac{1}{\epsilon}-\int_0^1 dt\log\frac{p^2t(1-t)+m^2}{\mu^2}). \tag{4.9}$$

We see that this divergence has the form of pole part proportional to $\frac{1}{\epsilon}$. To cancel it we must add to the initial kinetic term

$$S = \int d^8z\Phi(x,\theta)\bar{\Phi}(x,\theta) \tag{4.10}$$

(which is just $\int \frac{d^4p}{(2\pi)^4} d^4\theta \Phi(-p,\theta)\bar{\Phi}(p,\theta)$) a counterterm

$$\Delta S_{countr} = -\frac{\lambda^2}{32\pi^2\epsilon}\int d^8z\Phi(z)\bar{\Phi}(z) \tag{4.11}$$

which corresponds to the replacement of $\int d^8z\Phi\bar{\Phi}$ in the classical action by $\int d^8zZ\Phi(z)\bar{\Phi}(z)$ where

$$Z = 1 - \frac{\lambda^2}{32\pi^2\epsilon} \tag{4.12}$$

is a wave function renormalization.

The essential peculiarity of superfield theories is the fact that number of counterterms in these theories is less than in their non-supersymmetric analogs. For example, Wess-Zumino model is a supersymmetric generalization of ϕ^4-theory, but it possesses only renormalization of kinetic term and no renormalization of couplings. The conclusion about absence of divergent correction to coupling the λ (or as is the same – to chiral potential) is also called non-renormalization theorem. However, this theorem does not forbid **finite** corrections to superpotential which present in massless Wess-Zumino model [19, 20, 21, 22].

Theare are also some interesting properties of renormalization in superfield theories.

First, all tadpole-type contributions in Wess-Zumino model vanish: supergraph



has contribution proportional to $D^2\delta_{11} = \delta_{12}D^2\delta_{12} = 0$. The similar situation can occur in other superfield models involving the Wess-Zumino model as an ingredient. However, in theories including vertices proportional to integral over whole superspace (f.e. dilaton supergravity) tadpole contributions are not equal to zero [13].

Second, all contributions from vacuum supergraphs are proportional to $\int d^4\theta c$ (with c is a constant) and also vanish. However, this statement is not true for background dependent propagators. Using of background dependent propagators are very important method for calculation of effective action. Now we turn to its studying.

5 Effective action and loop expansion

Effective action is a central object of quantum field theory. Studying of effective action allows to investigate problems of vacuum stability, Green functions, spontaneous symmetry breaking, anomalies and many other problems.

Effective action in superfield theory is defined as usual as a generating functional of one-particle-irreducible Green functions. It is obtained as a Legendre transform for generating functional of connected Green functions:

$$\Gamma[\Phi] = W[J] - \int dzJ(z)\Phi(z). \tag{5.1}$$

Here $\Gamma[\Phi]$ is an effective action, dz denotes integral over the corresponding subspace (d^6z for chiral sources, d^8z for general ones), Φ is a set of all superfields, $\Phi(z)=\frac{\delta W[J]}{\delta J(x)}$ is so called mean field or background field, $W[J]=\frac{1}{i}\log Z[J]$ is a generating functional of the connected Green functions. As usual, $\Gamma[\Phi]$ satisfies the equation

$$\frac{\delta\Gamma[\Phi]}{\delta\Phi(x)}=-J(x).$$

The effective action can be expressed in the form of path integral [23]:

$$e^{\frac{i}{\hbar}\Gamma[\Phi]}=\int D\phi e^{\frac{i}{\hbar}(S[\phi]+\phi J-\Phi J)}. \tag{5.2}$$

Here $S[\phi]$ is a classical action of the corresponding theory. Note that ϕ is a variable of integration, and Φ is a function of classical source J which does not depend on ϕ. We introduced $\hbar$ by dimensional reasons and to obtain loop expansion. To calculate this integral we make change of variables of integration:

$$\phi\rightarrow\Phi+\sqrt{\hbar}\phi.$$

If we have several fields we can unite them into a column vector, and all consideration is quite analogous. The integral (5.2) after this change takes the form

$$e^{\frac{i}{\hbar}\Gamma[\Phi]}=\int D\phi e^{\frac{i}{\hbar}(S[\Phi+\sqrt{\hbar}\phi]+\sqrt{\hbar}\phi J)}. \tag{5.3}$$

Our aim consists here of the expansion of $\Gamma[\phi]$ in power series in $\hbar$ following the approach described in [23].

First, we expand factor in the exponent into power series in $\hbar$:

$$\begin{aligned}\frac{i}{\hbar}S[\Phi+\sqrt{\hbar}\phi]+\frac{i}{\sqrt{\hbar}}\phi J=\frac{i}{\hbar}\Big(S[\Phi]+S'[\Phi]\sqrt{\hbar}\phi+\frac{\hbar}{2}S''[\Phi]\phi^2+\ldots+\\ +\quad\frac{\hbar^{n/2}}{n!}S^{(n)}[\Phi]\phi^n+\ldots\Big).\end{aligned} \tag{5.4}$$

Here $S^{(n)}[\Phi]$ denotes n-th variational derivative of the classical action with respect to Φ (integration over corresponding space is assumed). This expansion can be substituted into (5.3). We introduce $\bar\Gamma[\Phi]=\Gamma[\Phi]-S[\Phi]$ which is a quantum contribution to effective action that can be expanded into power series in $\hbar$: $\bar\Gamma=\sum_{n=1}^{\infty}\hbar^n\Gamma^{(n)}$. As a result we have

$$e^{\frac{i}{\hbar}\bar\Gamma[\Phi]}=\int D\phi\exp\Big[\frac{i}{\hbar}\Big(S[\Phi]+S'[\Phi]\sqrt{\hbar}\phi+\frac{\hbar}{2}S''[\Phi]\phi^2+\ldots+\frac{\hbar^{n/2}}{n!}S^{(n)}[\Phi]\phi^n+\ldots\Big)\Big]. \tag{5.5}$$

Then, the block $\frac{i}{\sqrt{\hbar}}(S'[\Phi]+J)\phi$ can lead only to one-particle-reducible supergraphs since its contribution with one quantum field ϕ can form only one propagator. Hence we can omit this term. Then we can expand the exponent into power series in $\hbar$:

$$\begin{aligned}e^{\frac{i}{\hbar}\bar\Gamma[\Phi]}=\int D\phi e^{\frac{i}{2}S''[\Phi]\phi^2}\Big(1+\frac{i\sqrt{\hbar}}{3!}S^{(3)}[\Phi]\phi^3+\frac{i\hbar}{4!}S^{(4)}[\Phi]\phi^4+\\ +\quad(\frac{i\sqrt{\hbar}}{3!})^2(S^{(3)}[\Phi]\phi^3)^2+\ldots\Big).\end{aligned} \tag{5.6}$$

At the same time, after substituting the expansion of $\bar{\Gamma}$ in the left-hand side of (5.5) into power series in $\hbar$ we get $\exp(\frac{i}{\hbar}\bar{\Gamma}[\Phi]) = e^{i\Gamma^{(1)}[\Phi]}(1 + i\hbar\Gamma^{(2)}[\Phi] + \ldots)$ (here we suppose that $\hbar$ is a small parameter). Substituing this expansion into (5.6) and comparing equal powers of $\hbar$ we see that any correction $\Gamma^{(n)}$ corresponds to some correlator. For example, one-loop correction is defined from equation

$$\exp(i\Gamma^{(1)}[\Phi]) = \int D\phi e^{\frac{i}{2}S^{''}[\Phi]\phi^2}, \tag{5.7}$$

and two-loop one – from equation

$$\Gamma^{(2)} = \frac{1}{i}\frac{\int D\phi e^{(\frac{i}{2}S^{''}[\Phi]\phi^2)}\left(\frac{i}{4!}S^{(4)}[\Phi]\phi^4 - \frac{1}{(3!)2}(S^{(3)}[\Phi]\phi^3)^2\right)}{\int D\phi \exp(\frac{i}{2}S''[\Phi]\phi^2)}. \tag{5.8}$$

Here, as usual, integration over coordinates in expressions of the form $S^{(n)}[\Phi]\phi^n$ is assumed.

We can see that:

(i) All odd orders in $\sqrt{\hbar}$ vanish since they correspond to $\int D\phi\phi^{2n+1}\exp(\frac{i}{2}S^{''}[\Phi]\phi^2)$. Due to symmetrical properties this integral is equal to zero.

(ii) All terms beyond first order in $\hbar$ are expressed in the form of some correlators.

(iii) One-loop correction (5.7) can be expressed in the form of functional determinant since

$$\int D\phi \exp(\frac{i}{2}S^{''}[\Phi]\phi^2) = Det^{-1/2}S^{''}[\Phi], \tag{5.9}$$

which leads to

$$\Gamma^{(1)} = \frac{i}{2}Tr\log S^{''}[\Phi]. \tag{5.10}$$

And $S^{''}[\Phi]$ (further we denote it as Δ) is a some operator. In many cases it has the form $\Delta = \Box + \ldots$. We can express one-loop effective action in terms of functional (super)trace

$$\Gamma^{(1)} = \frac{i}{2}\mathrm{Tr}\int_0^\infty \frac{ds}{s}e^{is\Delta}. \tag{5.11}$$

This expression is called Schwinger representation for the one-loop effective action. Sign Tr denotes both matrix trace tr (if Δ possesses matrix indices) and functional trace, i.e.

$$\mathrm{Tr}e^{is\Delta} = \mathrm{tr}\int d^8z_1 d^8z_2 \delta^8(z_1 - z_2)e^{is\Delta}\delta^8(z_1 - z_2)$$

Calculation of $e^{is\Delta}$ in field theories is carried out with use of a special procedure called Schwinger-De Witt method or proper time method [24]. This method will be discussed in the next section.

Let us consider higher loop corrections. From (5.6) it is easy to see that all loop corrections beyond one-loop order have the form of some correlators, i.e. they include

$$\int D\phi \exp(\frac{i}{2}S^{''}[\Phi]\phi^2)\prod_n(S^{(n)}[\Phi]\phi^n). \tag{5.12}$$

Such correlators can be calculated in the way analogous to standard theory of perturbations. We can use expression

$$\int D\phi\phi^n e^{i\phi\Delta\phi} = (\frac{1}{i}\frac{\delta}{\delta j})^n \int D\phi e^{i(\phi\Delta\phi+j\phi)}|_{j=0}, \tag{5.13}$$

which allows to introduce diagram technique in which the role of vertices is played by $\frac{S^{(n)}[\Phi]\phi^n}{n!}$, and role of propagators – by Δ^{-1}. However since $\Delta = S''[\Phi]$ is background dependent (see above) we arrive at background dependent propagators $<\phi(z_1)\phi(z_2)>= \Delta^{-1}\delta^8(z_1-z_2)$. These propagators are known to be found exactly only in some special cases, the most important of them are: first, constant in space-time background superfields, second, the background superfields are only chiral. Further we consider some examples.

Let us turn again to (5.6). We see that each quantum superfield corresponds to $\hbar^{-1/2}$, and each vertex – to $\hbar^{-1}$ (which provides $\hbar^{n/2-1}S^{(n)}[\Phi]\phi^n$). Arbitrary (super)graph with P propagators and V vertices contain $2P$ quantum superfields (each propagator is formed by contraction of two superfields). Therefore if this (super)graph contain vertices $S^{n_1}[\Phi]\phi^{n_1}, S^{n_2}[\Phi]\phi^{n_2}, \dots, S^{n_V}[\Phi]\phi^{n_V}$ its power in $\hbar$ is $\sum_{i=1}^V(\frac{n_i}{2}-1) = \frac{1}{2}\sum_{i=1}^V n_i - V$. However, $\sum_{i=1}^V n_i$ is just the number of quantum fields associated with all vertices which is equal to $2P$. Therefore the correlator described by this (super)graph has power of $\hbar$ equal to $P-V=L-1$, with L is number of loops. But any correlator of the form (5.6) is a contrbution to $\frac{\Gamma}{\hbar}$, hence contribution from L-loop (super)graph to Γ is proportional to $\hbar^L$. Hence we found that the order in $\hbar$ from an arbitrary (super)graph is just the number of loops in it, and the expansion in powers of $\hbar$ is called loop expansion. As a result we see that loop corrections can be calculated on the base of special (super)field technique.

Let us make some comments. One of the most frequent questions is: how is the definition of (one-loop) correction in effective action in terms of trace of logarithm related to expression of the same correction in terms of supergraphs?

To clarify this relation we give an example. One-loop effective action in Wess-Zumino model is given by [25, 10]

$$\Gamma^{(1)} = \frac{i}{2}\mathrm{Tr}\log(\Box - \frac{1}{4}\Psi\bar{D}^2 - \frac{1}{4}\bar{\Psi}D^2). \tag{5.14}$$

Here Ψ is background chiral superfield. This expression can be rewritten as

$$\Gamma^{(1)} = \frac{i}{2}\mathrm{Tr}\log[\Box(1 - \frac{1}{4\Box}(\Psi\bar{D}^2 + \bar{\Psi}D^2))]. \tag{5.15}$$

Expansion of the logarithm into power series leads to

$$\Gamma^{(1)} = \frac{i}{2}\mathrm{Tr}\sum_{n=1}^{\infty}\frac{1}{n}[\frac{1}{4\Box}(\Psi\bar{D}^2 + \bar{\Psi}D^2)]^n. \tag{5.16}$$

This expression exactly reproduces the total contribution for the sum of the following supergraphs



Fig.4

External lines here are for alternating $\Psi\bar{D}^2/4$ and $\bar{\Psi}D^2/4$, and internal ones are for $\Box^{-1}$. At the same time, if we consider theory of real scalar superfield u in external chiral superfield Ψ with action

$$S=\int d^8zu(\Box-\frac{1}{4}\Psi\bar{D}^2-\frac{1}{4}\bar{\Psi}D^2)u, \tag{5.17}$$

it leads just to these supergraphs (if $\int d^8zu(-\frac{1}{4}\Psi\bar{D}^2)u$ and the conjugated term are treated as vertices), and one-loop effective action for this theory is again given by (5.14).

We can see that the expression of one-loop effective action in the form of the trace of the logarithm of some operator allows to use some special technique which is equivalent to supergraph approach, but more convenient in many cases. This technique is called proper-time technique.

6 Superfield proper-time technique

As we have already proved, if the quadratic action of a quantum (super)field ϕ on classical background Φ has the form $\int dx\phi\Delta[\Phi]\phi$ ($\int dx$ here denotes integral over all (super)space), one-loop effective action in this theory is $\Gamma^{(1)}=\frac{i}{2}Tr\int_0^\infty\frac{ds}{s}e^{is\Delta}$. Therefore we face the problem of calculating the operator $e^{is\Delta}$. In most important cases $\Delta=\Box+\dots$ where dots denote background dependent terms. It is known [24] that the best way to find this operator in the case of common field theory is as follows. We introduce $U(x,x'|s)=e^{is\Delta}\delta^4(x-x')$ called Schwinger kernel. Of course, U depends on background superfields. It satisfies the equation:

$$i\frac{\partial U}{\partial s}=-U\Delta. \tag{6.1}$$

The Δ is supposed to have form of power series in derivatives. And U satisfies initial condition

$$U(x,x')|_{s=0}=\delta^4(x-x').$$

In general case U is represented in the form of infinite power series in parameter s (called proper time) as [24]

$$U=-\frac{i}{(4\pi s)^2}\exp(\frac{i}{4s}(x-x')^2)\sum_{n=0}^{\infty}a_n(is)^n. \tag{6.2}$$

(Ultraviolet) divergences correspond to **lower** orders of this expansion (note that ultraviolet case corresponds to $s\to 0$, infrared one – to $s\to\infty$). Coefficients a_n depend on background superfields and their derivatives. We note that if background superfields are put to zero, we arrive at

$$U^{(0)}(x,x';s)=e^{is\Box}\delta^4(x-x')=-\frac{i}{(4\pi s)^2}\exp(\frac{i}{4s}(x-x')^2), \tag{6.3}$$

which satisfies condition

$$i\int_0^\infty dsU^{(0)}(x,x';s)=\frac{1}{\Box}\delta^4(x-x'). \tag{6.4}$$

The approach in case of superfield theories is quite analogous. However, it possesses an essential advantage. In this case it is more convenient to expand Schwinger kernel $U(x,x';s)$ not in infinite power series in s but in a **finite** power series in spinor supervcovariant derivatives (these series are finite due to anticommutation properties of spinor derivatives).

Really, in most cases operator Δ in superfield theories looks like

$$\Delta=\Box+\sum A_{nm}(D^\alpha)^n(\bar{D}^{\dot\alpha})^m\equiv\Box+\tilde{\Delta} \tag{6.5}$$

with $\tilde{\Delta}$ is some background dependent operator (in most cases it contains only even orders in spinor derivatives, here we consider this case), A_{nm} are background dependent coefficients. We introduce the structure

$$U(z,z';s)=\exp(is\Delta)\delta^8(z-z')\equiv\exp(is\tilde{\Delta})\exp(is\Box)\delta^8(z-z') \tag{6.6}$$

(last identity is valid in the case of contributions which do not depend on space-time derivatives of superfields). We substitute natural initial condition

$$U(z,z';s)|_{s=0}=\delta^8(z-z').$$

And $\exp(is\Box)\delta^8(z-z')=\delta^4(\theta-\theta')U^{(0)}(x,x';s)$ where $U^{(0)}(x,x';s)$ is given by (6.3). Hence

$$U(z,z';s)=\exp(is\tilde{\Delta})U^{(0)}(x,x';s)\delta^4(\theta-\theta'). \tag{6.7}$$

Therefore we face the problem of calculating $\tilde{U}=\exp(is\tilde{\Delta})$. The $\tilde{U}$ satisfies the equation

$$i\frac{\partial\tilde{U}}{\partial s}=-\tilde{U}\tilde{\Delta}. \tag{6.8}$$

It is easy to see that $\tilde{U}|_{s=0}=1$. We expand $\tilde{U}$ into power series in spinor supercovariant derivatives:

$$\begin{aligned}\tilde{U} &= 1+\frac{1}{16}A(s)D^2\bar{D}^2+\frac{1}{16}\tilde{A}(s)\bar{D}^2D^2+\frac{1}{8}B^\alpha(s)D_\alpha\bar{D}^2+\frac{1}{8}\tilde{B}_{\dot\alpha}\bar{D}^{\dot\alpha}D^2+\\ &+ \frac{1}{4}C(s)D^2+\frac{1}{4}\tilde{C}(s)\bar{D}^2\end{aligned} \tag{6.9}$$

and substitute (6.9) into equation (6.8). As a result we obtain in right-hand side some power series in spinor derivatives. Comparing coefficients at analogous derivatives in right-hand side and left-hand side of identity we get

$$\begin{aligned}
\frac{1}{16}\dot{A} &= \tilde{U}\tilde{\Delta}|_{D^2\bar{D}^2};\\
\frac{1}{8}\dot{B}^{\alpha} &= \tilde{U}\tilde{\Delta}|_{D_\alpha\bar{D}^2};\\
\frac{1}{4}\dot{C} &= \tilde{U}\tilde{\Delta}|_{D^2} \qquad (6.10)
\end{aligned}$$

and analogous equations for $\tilde{A}, \tilde{B}_{\dot{\alpha}}, \tilde{C}$. Here dot denotes $\frac{\partial}{\partial is}$, and $|_{D^2}$ etc. denotes coefficient at D^2 etc. in $\tilde{U}\tilde{\Delta}$. As a result we have system of first-order differential equations on coefficients determining structure of operator $\tilde{U}$. Since $\tilde{U}|_{s=0}=1$ we have natural initial conditions

$$A|_{s=0}=\tilde{A}|_{s=0}=B^{\alpha}|_{s=0}=\tilde{B}_{\dot{\alpha}}|_{s=0}=C|_{s=0}=\tilde{C}|_{s=0}=0. \qquad (6.11)$$

The system (6.10) with initial conditions (6.11) can be solved like common system of differential equations (note however, that this solution is mostly found in special cases, such as independence on spinor derivatives of background superfields, or dependence on chiral background superfields only etc.)

Then, $\tilde{U}(s)$ (often called heat kernel) can be used for calculation of Green function as

$$G(z_1,z_2)=i\int_0^\infty ds\tilde{U}U^{(0)}(x,x';s)\delta^4(\theta-\theta') \qquad (6.12)$$

(note that $\tilde{U}$ is a differential operator in superspace) and for calculation of one-loop effective action as

$$\Gamma^{(1)}=\frac{i}{2}\int_0^\infty\frac{ds}{s}\int d^8zd^8z'\delta^8(z-z')\tilde{U}U^{(0)}(x,x';s)\delta^4(\theta-\theta'). \qquad (6.13)$$

As usual, $\int d^8z=\int d^4xd^4\theta$, we also use definition (6.3). Then, it is known that $\delta^4(\theta-\bar{\theta})D^2\bar{D}^2\delta^4(\theta-\bar{\theta})=16\delta^4(\theta-\bar{\theta})$, and all products of less number of spinor derivatives give zero trace. Hence only coefficients of (6.9) giving non-zero contribution to one-loop effective action are A and $\tilde{A}$. And one-loop effective action looks like

$$\Gamma^{(1)}=\frac{i}{2}\int_0^\infty\frac{ds}{s}\int d^4xd^4\theta(A(s)+\tilde{A}(s))U^{(0)}(x,x';s)|_{x=x'}. \qquad (6.14)$$

As a result we developed technique for calculating background dependent propagators and one-loop effective action. Application of this technique will be further considered on examples of several theories. There is an essential modification of this method for supergauge theories [26].

7 Problem of superfield effective potential

Effective potential in standard quantum field theory is defined as the effective Lagrangian considered at constant values of scalar fields, and other fields are put to zero. The effective potential is used for studying of spontaneous symmetry breaking and vacuum stability [27].

First, let us shortly describe effective potential in common quantum field theory. The effective action has the form

$$\Gamma[\phi]=\int d^4x(-V_{eff}(\phi)-\frac{1}{2}Z(\phi)\partial_m\phi\partial^m\phi+\ldots), \tag{7.1}$$

where $Z(\phi)$ is a some function of ϕ, and $V_{eff}(\phi)$ is effective potential. For slowly varying fields, therefore,

$$\Gamma[\phi]=-\int d^4xV_{eff}(\phi),$$

therefore effective potential is a low-energy leading term. It can be represented in the form of loop expansion

$$V_{eff}(\phi)=V(\phi)+\sum_{n=1}^{\infty}\hbar^nV^{(n)}(\phi). \tag{7.2}$$

For example, consider the theory with action

$$S=\int d^4x(\frac{1}{2}\phi\Box\phi+V(\phi)). \tag{7.3}$$

After background-quantum splitting $\phi\to\Phi+\chi$ where Φ is background superfield and χ is quantum one, we find the quadratic action of quantum superfields

$$S_2=\int d^4x[\frac{1}{2}\chi(\Box+V^{''}(\Phi))\chi] \tag{7.4}$$

which leads to one-loop effective action $\Gamma^{(1)}[\Phi]$ of the form

$$\Gamma^{(1)}[\Phi]=\frac{i}{2}Tr\log(\Box+V^{''}(\Phi)). \tag{7.5}$$

Following Section 5, we can express this trace of logarithm in the form of diagrams:



Fig. 5

where external lines are $V^{''}[\Phi]$. Internal lines correspond to $\frac{1}{\Box}\delta^4(\theta_1-\theta_2)$.

The sum of contributions from these supergraphs is

$$S=\sum_{n=1}^{\infty}\frac{1}{n}\int\frac{d^4k}{(2\pi)^4}(-\frac{V^{''}(\Phi)}{k^2})^n=-\int\frac{d^4k}{(2\pi)^4}\log(1+\frac{V^{''}(\Phi)}{k^2}), \tag{7.6}$$

which after integration over d^4k and extraction of divergences is equal to [10]

$$\frac{1}{64\pi^2}(V''(\Phi))^2(\log\frac{V''(\Phi)}{\mu^2}+C), \tag{7.7}$$

where C is some constant. The same result can be obtained via proper-time method (see calculation f.e. in [10]).

Now we turn to superfield case. Let $\Gamma[\Phi,\bar{\Phi}]$ be the renormalized effective action for a theory of chiral and antichiral superfields. We can represent it as

$$\begin{aligned}\Gamma[\bar{\Phi},\Phi] &= \int d^8z\mathcal{L}_{eff}(\Phi,D_A\Phi,D_AD_B\Phi;\bar{\Phi},D_A\bar{\Phi},D_AD_B\bar{\Phi})+\\ &+ (\int d^6z\mathcal{L}^{(c)}_{eff}(\Phi)+h.c.)+\ldots \end{aligned}\tag{7.8}$$

Here $D_A\Phi, D_AD_B\Phi,\ldots$ are possible supercovariant derivatives of superfields $\Phi,\bar{\Phi}$. The term $\mathcal{L}_{eff}$ is called general effective Lagrangian, and $\mathcal{L}^{(c)}_{eff}$ is called chiral effective Lagrangian. Both these effective Lagrangians can be expanded into power series in supercovariant derivatives of background superfields. Dots denote terms depending on space-time derivatives of Φ, $\bar{\Phi}$. We note that since chiral effective Lagrangian by definition depends only on Φ but not on $\bar{D}^2\bar{\Phi}$ all terms of the form

$$\int d^6z\Phi^n(\bar{D}^2\bar{\Phi})^m$$

using relation $\int d^6z(-\frac{\bar{D}^2}{4})=\int d^8z$ can be rewritten as

$$\int d^8z\Phi^n\bar{\Phi}(\bar{D}^2\bar{\Phi})^{m-1},$$

i.e. in the form corresponding to general effective Lagrangian. Therefore here and further we consider all formally chiral expressions but involving $(\bar{D}^2\bar{\Phi})^m$ as contributions to general effective Lagrangian.

We note that all chiral contributions can be also represented as integral over whole superspace:

$$\int d^6zG(\Phi)=\int d^8z(-\frac{D^2}{4\Box})G. \tag{7.9}$$

Further, in component approach we must put scalar component fields to constants, and spinor ones – to zero, f.e. in Wess-Zumino model we write

$$A=const,\ F=const,\ \psi_\alpha=0.$$

However, this condition is not supersymmetric, therefore we use condition of superfield **constant in space-time**:

$$\partial_a\Phi=0. \tag{7.10}$$

Since ∂_a commutes with all generators of supersymmetry, this condition is supersymmetric.

Effective potential is introduced as

$$V_{eff} = \Big\{ -\int d^4\theta \mathcal{L}_{eff} - (\int d^2\theta \mathcal{L}^{(c)}_{eff} + h.c.)\Big\}|_{\partial_a\Phi=\partial_a\bar{\Phi}=0}. \tag{7.11}$$

The minus sign is put by convention. We can introduce general effective potential $\mathcal{L}_{eff}|_{\partial_a\Phi=\partial_a\bar{\Phi}=0}$ and chiral effective potential $\mathcal{L}^{(c)}_{eff}|_{\partial_a\Phi=0}$. It is easy to see that the general effective potential can be expressed as

$$\mathcal{L}_{eff} = \mathbf{K}(\Phi,\bar{\Phi}) + \mathbf{F}(D_\alpha\Phi, \bar{D}_{\dot{\alpha}}\bar{\Phi}, D^2\Phi, \bar{D}^2\bar{\Phi}; \Phi, \bar{\Phi}) \tag{7.12}$$

with $\mathbf{F}|_{D_\alpha\Phi,\bar{D}_{\dot{\alpha}}\bar{\Phi},D^2\Phi,\bar{D}^2\bar{\Phi}=0} = 0$. The $\mathbf{K}$ is called kählerian effective potential, and $\mathbf{F}$ is called auxiliary fields' effective potential, it is at least of third order in auxiliary fields of Φ and $\bar{\Phi}$. These objects can be represented in the form of loop expansion:

$$\mathbf{K}(\Phi,\bar{\Phi}) = K_0(\Phi,\bar{\Phi}) + \sum_{L=1}^{\infty} \hbar^L K_L(\Phi,\bar{\Phi}) \tag{7.13}$$

$$\mathbf{F} = \sum_{L=1}^{\infty} \hbar^L F_L \tag{7.14}$$

(the term corresponding to $L=0$ in the expression for $\mathbf{F}$ is absent for theories which do not include derivative depending terms in the classical action, such as the Wess-Zumino model), and

$$\mathcal{L}^{(c)}_{eff}(\Phi) = \mathcal{L}^{(c)}(\Phi) + \sum_{L=1}^{\infty} \hbar^L \mathcal{L}^{(c)}_L(\Phi). \tag{7.15}$$

Here $K_L, F_L, \mathcal{L}^{(c)}_L$ are quantum corrections. For Wess-Zumino model $\mathcal{L}^{(c)}_1 = 0$, however, in some quantum theories (f.e. in $N=1$ super-Yang-Mills theory with chiral matter) one-loop contribution to chiral effective potential exists [20].

The expansion of the effective potential by the rules (7.11–7.15) can be applied for all superfield theories including noncommutative ones. However, we note that effective potential in theories including gauge superfields must depend on them in a special way. Really, effective action in such theories should be expressed in terms of some gauge convariant constructions, f.e. in background field method gauge superfield is either incorporated to chiral superfields or presents in supercovariant derivatives and gauge invariant superfield strengths [28, 10].

Let us give a few remarks about the method of calculating effective potential. The best way for it is, of course, using of background dependent propagators which are expressed in terms of common propagators and background superfields. Background dependent propagators can be in certain cases exactly found. To calculate kählerian effective potential and auxiliary fields' effective potential one can straightforwardly omit all space-time derivatives, moreover, to study kählerian effective potential one can omit **ALL** supercovariant derivatives and treate background superfields as constants until final integration. The calculation of chiral effective potential, however, is characterized by some difficulties. We will study an approach to it on example of Wess-Zumino model.

8 Wess-Zumino model and problem of chiral effective potential

Now we turn to consideration of superfield effective potential in Wess-Zumino model. Here we follow the papers [21, 22, 25] and book [10].

The superfield action of Wess-Zumino model is given by (2.9). Following loop expansion approach we carry out background-quantum splitting by the rule

$$\begin{aligned} \Phi &\to \Phi + \sqrt{\hbar}\phi; \\ \bar{\Phi} &\to \bar{\Phi} + \sqrt{\hbar}\bar{\phi}. \end{aligned} \tag{8.1}$$

The expression (5.6) defining effective action under such changes takes the form (here $\bar{\Gamma} = \sum_{L=1}^{\infty} \hbar^L \Gamma_L$)

$$\begin{aligned} e^{\frac{i}{\hbar}\bar{\Gamma}[\Phi,\bar{\Phi}]} &= \int D\phi D\bar{\phi} \exp\Big(\frac{i}{2}\begin{pmatrix} \phi\bar{\phi} \end{pmatrix}\begin{pmatrix} \psi & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & \psi \end{pmatrix}\begin{pmatrix} \phi \\ \bar{\phi} \end{pmatrix} + \\ &+ \frac{i\sqrt{\hbar}}{3!}(\frac{\lambda}{3!}\phi^3 + h.c.)\Big). \end{aligned} \tag{8.2}$$

The quadratic action of quantum superfields looks like

$$S^{(2)} = \frac{1}{2}\begin{pmatrix} \phi\bar{\phi} \end{pmatrix}\begin{pmatrix} \psi & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & \psi \end{pmatrix}\begin{pmatrix} \phi \\ \bar{\phi} \end{pmatrix}. \tag{8.3}$$

And the matrix superpropagator by definition is an operator inverse to

$$\begin{pmatrix} \psi & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & \psi \end{pmatrix}\begin{pmatrix} \delta_+ & 0 \\ 0 & \delta_- \end{pmatrix}. \tag{8.4}$$

We can see that this matrix superpropagator can be represented in the form

$$G(z_1, z_2) = \begin{pmatrix} G_{++}(z_1, z_2) & G_{+-}(z_1, z_2) \\ G_{-+}(z_1, z_2) & G_{--}(z_1, z_2) \end{pmatrix}. \tag{8.5}$$

where + denotes chirality with respect to corresponding argument, and − correspondingly – antichirality.

It turns out to be that in Wess-Zumino model this matrix looks like

$$G(z_1, z_2) = \frac{1}{16}\begin{pmatrix} \bar{D}_1^2\bar{D}_2^2 G_v^{\psi}(z_1, z_2) & \bar{D}_1^2 D_2^2 G_v^{\psi}(z_1, z_2) \\ D_1^2\bar{D}_2^2 G_v^{\psi}(z_1, z_2) & D_1^2 D_2^2 G_v^{\psi}(z_1, z_2) \end{pmatrix}. \tag{8.6}$$

where $G_v^{\psi}(z_1, z_2) = (\Box + \frac{1}{4}\psi\bar{D}^2 + \frac{1}{4}\bar{\psi}D^2)^{-1}\delta^8(z_1 - z_2)$. Really, consider relation

$$\begin{pmatrix} \psi & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & \psi \end{pmatrix}\frac{1}{16}\begin{pmatrix} \bar{D}_1^2\bar{D}_2^2 G_v^{\psi}(z_1, z_2) & \bar{D}_1^2 D_2^2 G_v^{\psi}(z_1, z_2) \\ D_1^2\bar{D}_2^2 G_v^{\psi}(z_1, z_2) & D_1^2 D_2^2 G_v^{\psi}(z_1, z_2) \end{pmatrix} = -\begin{pmatrix} \delta_+ & 0 \\ 0 & \delta_- \end{pmatrix} \tag{8.7}$$

and act on both parts of this relation with the operator

$$\begin{pmatrix} 0 & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & 0 \end{pmatrix}.$$

We get the following system of equations on components of matrix superpropagator G:

$$\begin{aligned}
\Box G_{++} &- \frac{1}{4}\bar{D}_1^2(\bar{\psi}G_{-+}) = 0; \\
\Box G_{-+} &- \frac{1}{4}D_1^2\psi G_{++}) = \frac{1}{16}D_1^2\bar{D}_2^2\delta^8(z_1 - z_2); \\
\Box G_{--} &- \frac{1}{4}D_1^2(\psi G_{+-}) = 0; \\
\Box G_{+-} &- \frac{1}{4}\bar{D}_1^2(\bar{\psi}G_{--}) = \frac{1}{16}\bar{D}_1^2 D_2^2\delta^8(z_1 - z_2).
\end{aligned} \tag{8.8}$$

Straightforward checking shows that components $G_{++}, G_{+-}, G_{-+}, G_{--}$ given by (8.6) satisfy this equation. Thus, we found matrix superpropagator (8.6) which will be used for calculation of loop corrections.

Let us consider the one-loop effective action. Formally it has the form

$$\Gamma^{(1)} = -\frac{i}{2}\mathrm{Tr}\log G$$

where matrix superpropagator G is given by (8.6). However, straightforward calculation of this trace is very complicated since the elements of this matrix are defined in different subspaces. The one-loop effective action $\Gamma^{(1)}$ can be obtained from relation

$$e^{i\Gamma^{(1)}} = \int D\phi D\bar{\phi}\exp\Big(\frac{i}{2}\begin{pmatrix} \phi\bar{\phi} \end{pmatrix}\begin{pmatrix} \psi & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & \psi \end{pmatrix}\begin{pmatrix} \phi \\ \bar{\phi} \end{pmatrix}\Big). \tag{8.9}$$

To take this integral we introduce a trick [25] which is used in many theories describing dynamics of chiral superfields.

We consider theory of real scalar superfield with action

$$S = \frac{1}{16}\int d^8 z v D^\alpha \bar{D}^2 D_\alpha v. \tag{8.10}$$

The action is invariant under gauge transformations $\delta v = \Lambda - \bar{\Lambda}$ (here Λ is chiral, and $\bar{\Lambda}$ is antichiral). According to Faddeev-Popov approach, the effective action W for this theory can be introduced as

$$e^{iW} = \int Dv e^{\frac{i}{16}\int d^8 z v D^\alpha \bar{D}^2 D_\alpha v}\delta(\chi). \tag{8.11}$$

Here $\delta(\chi)$ is a functional delta function, and χ is a gauge-fixing function. We choose χ in the form of column matrix

$$\chi = \begin{pmatrix} \frac{1}{4}D^2 v - \bar{\phi} \\ \frac{1}{4}\bar{D}^2 v - \phi \end{pmatrix}.$$

Note that since supercovariant derivatives are not real we must impose two conditions, and (8.11) takes the form

$$e^{iW}=\int Dv e^{\frac{i}{16}\int d^8z vD^{\alpha}\bar{D}^2D_{\alpha}v}\delta(\frac{1}{4}D^2v-\bar{\phi})\delta(\frac{1}{4}\bar{D}^2v-\phi)\mathrm{det}\Delta, \tag{8.12}$$

where

$$\Delta=\left(\begin{array}{cc} -\frac{1}{4}\bar{D}^2 & 0 \\ 0 & -\frac{1}{4}D^2 \end{array}\right)$$

is a Faddeev-Popov matrix. We note that W is constant by construction. We multiply left-hand sides and right-hand sides of (8.9) and (8.12) respectively, as a result we arrive at

$$\begin{aligned} e^{i\Gamma^{(1)}+W} &= \int D\phi D\bar{\phi}Dv \exp\Big(\frac{i}{2}\left(\begin{array}{c} \phi\bar{\phi} \end{array}\right)\left(\begin{array}{cc} \psi & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & \psi \end{array}\right)\left(\begin{array}{c} \phi \\ \bar{\phi} \end{array}\right)+\frac{i}{16}vD^{\alpha}\bar{D}^2D_{\alpha}v\Big)\times \\ &\times \delta(\frac{1}{4}D^2v-\bar{\phi})\delta(\frac{1}{4}\bar{D}^2v-\phi)\mathrm{det}\Delta. \end{aligned} \tag{8.13}$$

Integration over $\phi,\bar{\phi}$ with use of delta functions leads to

$$e^{i\Gamma^{(1)}+W} = \int D\phi D\bar{\phi}\exp\Big(\frac{i}{2}v(\Box-\frac{1}{4}\psi\bar{D}^2-\frac{1}{4}\bar{\psi}D^2)v\Big)\mathrm{det}\Delta. \tag{8.14}$$

However, W and $\mathrm{det}\Delta$ are constants which can be omitted. We also took into account that $\frac{1}{16}\{D^2,\bar{D}^2\}+\frac{1}{8}D^{\alpha}\bar{D}^2D_{\alpha}=\Box$ [15], hence the one-loop effective action is equal to

$$\Gamma^{(1)}=\frac{i}{2}\mathrm{Tr}\log(\Box-\frac{1}{4}\psi\bar{D}^2-\frac{1}{4}\bar{\psi}D^2) \tag{8.15}$$

Here as usual $\psi=m+\lambda\Phi,\bar{\psi}=m+\lambda\bar{\Phi}$. This one-loop effective action can be expressed in form of Schwinger expansion:

$$\Gamma^{(1)}=\frac{i}{2}\mathrm{Tr}\int_0^{\infty}\frac{ds}{s}\exp(\Box-\frac{1}{4}\psi\bar{D}^2-\frac{1}{4}\bar{\psi}D^2), \tag{8.16}$$

or, after manifest writing the trace,

$$\Gamma^{(1)}=\frac{i}{2}\int d^8z_1d^8z_2\int_0^{\infty}\frac{ds}{s}\delta^8(z_1-z_2)\exp(is(-\frac{1}{4}\psi\bar{D}^2-\frac{1}{4}\bar{\psi}D^2))e^{is\Box}\delta^8(z_1-z_2). \tag{8.17}$$

We consider kernel $\tilde{U}(\psi|s)=\exp(is(-\frac{1}{4}\psi\bar{D}^2-\frac{1}{4}\bar{\psi}D^2))\equiv e^{is\Delta}$. It evidently satisfies the equation

$$\frac{\partial\tilde{U}}{\partial s}=i\tilde{U}\Delta.$$

It turns out to be that if we calculate kählerian effective potential and all supercovariant derivatives from background superfields $\psi,\bar{\psi}$ are omitted this equation can be easily solved. We express $\tilde{U}$ in the form (6.9). Then $\tilde{U}\Delta$ is equal to

$$\begin{aligned}
\tilde{U}\Delta &= -\frac{1}{4}\psi\bar{D}^2 - \frac{1}{4}\bar{\psi}D^2 - \\
&- \frac{1}{4}\psi\tilde{A}\Box\bar{D}^2 - \frac{1}{4}\bar{\psi}A\Box D^2 + \\
&+ \frac{1}{4}\tilde{B}_{\dot{\alpha}}\psi\partial^{\alpha\dot{\alpha}}D_\alpha\bar{D}^2 - \frac{1}{4}B_\alpha\bar{\psi}\partial^{\alpha\dot{\alpha}}\bar{D}_{\dot{\alpha}}D^2 - \\
&- \frac{1}{16}\bar{\psi}\bar{C}\bar{D}^2D^2 - \frac{1}{16}\psi C D^2\bar{D}^2. \qquad (8.18)
\end{aligned}$$

Comparing coefficients at analogous derivatives in $\frac{1}{i}\frac{\partial\tilde{U}}{\partial s}$ and $\tilde{U}\Delta$ we get the following system of equations

$$\begin{aligned}
\dot{A} &= -\psi C; \\
\dot{B}^\alpha &= 2i\tilde{B}_{\dot{\alpha}}\psi\partial^{\alpha\dot{\alpha}}; \\
\dot{C} &= -\bar{\psi} - \bar{\psi}A\Box. \qquad (8.19)
\end{aligned}$$

System for $\tilde{A}, \tilde{B}, \tilde{C}$ has the analogous form with changing $\psi \to \bar{\psi}, A \to \tilde{A}$ etc. Here dot denotes $\frac{1}{i}\frac{\partial}{\partial s} \equiv \frac{\partial}{\partial\tilde{s}}$. Since $\tilde{U}|_{s=0} = 1$, and all terms in expansion of $\tilde{U}$ (6.9) are evidently linearly independent, natural initial conditions are

$$A = \tilde{A} = B^\alpha = \tilde{B}_{\dot{\alpha}} = C = \tilde{C}|_{s=0} = 0. \qquad (8.20)$$

We find that the system of equations for B^α and $\tilde{B}_{\dot{\alpha}}$ is closed (it is separated from whole system (8.19)) and homogeneous. Initial conditions above make its only solution to be zero, $B^\alpha, \tilde{B}_{\dot{\alpha}} = 0$. The remaining from (8.19) system for A and C (and analogous one for $\tilde{A}$ and $\tilde{C}$) can be easily solved like standard system of common first-order differential equations. Its solution looks like

$$\begin{aligned}
C &= -\sqrt{\frac{\bar{\psi}\Box}{\psi}}(A_0^1\exp(i\omega s) - A_0^2\exp(-i\omega s)) \\
A &= A_0^1\exp(i\omega s) + A_0^2\exp(-i\omega s) - \frac{1}{\Box} \qquad (8.21)
\end{aligned}$$

Here $\omega = \sqrt{\psi\bar{\psi}\Box}$. Imposing initial conditions (8.20) allows to fix coefficients A_0^1, A_0^2. As a result we get

$$\begin{aligned}
C &= -\sqrt{\frac{\bar{\psi}}{\psi\Box}}\sinh(is\sqrt{\psi\bar{\psi}\Box}) \\
A &= \frac{1}{\Box}[\cosh(is\sqrt{\psi\bar{\psi}\Box}) - 1] \qquad (8.22)
\end{aligned}$$

Since A is symmetric with respect to change $\psi \to \bar{\psi}$ we find that $A = \tilde{A}$. We note that only A and $\tilde{A}$ contribute to trace in (8.17). Therefore one-loop kählerian contribution to effective action is equal to

$$K^{(1)} = \frac{i}{2}\int d^4x d^4\theta \int_0^\infty \frac{d\tilde{s}}{\tilde{s}}\frac{1}{\Box}[\cosh(\tilde{s}\sqrt{\psi\bar{\psi}\Box}) - 1]U_0(x, x'; s)|_{x=x'}. \qquad (8.23)$$

Here $U_0(x,x';s)$ is given by (6.3). This function satisfies the equation (see section 6):

$$\Box^n U_0(x,x';s)|_{x=x'} = (\frac{\partial}{\partial\tilde{s}})^n \frac{-i}{16\pi^2\tilde{s}^2}.$$

We expand (8.23) into power series:

$$\frac{1}{\Box}[\cosh(\tilde{s}\sqrt{\psi\bar{\psi}\Box}) - 1] = \sum_{n=0}^{\infty} \tilde{s}^{2n+2}\frac{(\psi\bar{\psi})^{n+1}}{(2n+2)!}\Box^n.$$

And

$$\begin{aligned} K^{(1)} &= \frac{i}{2}\int d^4x d^4\theta \int_0^\infty \frac{d\tilde{s}}{\tilde{s}}\sum_{n=0}^{\infty}\tilde{s}^{2n+2}\frac{(\psi\bar{\psi})^{n+1}}{(2n+2)!}\Box^n U_0(x,x';s)|_{x=x'} = \\ &= -\frac{i}{2}\int d^4x d^4\theta \int_0^\infty \frac{d\tilde{s}}{\tilde{s}}\sum_{n=0}^{\infty}\tilde{s}^{2n+2}\frac{(\psi\bar{\psi})^{n+1}}{(2n+2)!}(\frac{\partial}{\partial\tilde{s}})^n\frac{-i}{16\pi^2\tilde{s}^2} = \\ &= -\frac{1}{32\pi^2}\int d^8z \int_{L^2}^\infty \frac{d\tilde{s}}{\tilde{s}^2}\sum_{n=0}^{\infty}(-1)^n\frac{(\tilde{s}\psi\bar{\psi})^{n+1}(n+1)!}{(2n+2)!}. \end{aligned} \quad (8.24)$$

Here we cut integral at lower limit by introducing L^2 for regularization. We make the change $\tilde{s}\psi\bar{\psi} = t$. As a result, the one-loop kählerian contribution to effective action takes the form

$$K^{(1)} = -\frac{1}{32\pi^2}\int d^8z\psi\bar{\psi}\int_{\psi\bar{\psi}L^2}^\infty dt\sum_{n=0}^{\infty}\frac{(n+1)!t^{n+1}(-1)^n}{(2n+2)!} \quad (8.25)$$

Then, $\sum_{n=0}^{\infty}\frac{(n+1)!t^{n+1}(-1)^n}{(2n+2)!} = t\int_0^1 due^{-\frac{t}{4}(1-u^2)}$. Hence

$$K^{(1)} = -\frac{1}{32\pi^2}\int d^8z\psi\bar{\psi}\int_{\psi\bar{\psi}L^2}^\infty \frac{dt}{t}\int_0^1 due^{-\frac{t}{4}(1-u^2)}. \quad (8.26)$$

At $L^2 \to 0$ this integral tends to

$$K^{(1)} = -\frac{1}{32\pi^2}\psi\bar{\psi}\log(\mu^2L^2) - \frac{1}{32\pi^2}\psi\bar{\psi}(\log\frac{\psi\bar{\psi}}{\mu^2} - \xi). \quad (8.27)$$

where ξ is some constant which can be absorbed into redefinition of μ. We can add the counterterm $\frac{1}{32\pi^2}\psi\bar{\psi}\log(\mu^2L^2)$ to cancel the divergence. Such a counterterm corresponds to renormalization of kinetic term by the rule

$$\Phi \to Z^{1/2}\Phi;\ Z = 1 + \frac{\lambda^2}{32\pi^2}\log(\mu^2L^2). \quad (8.28)$$

And the renormalized kählerian effective potential is

$$K^{(1)} = -\frac{1}{32\pi^2}\psi\bar{\psi}(\log\frac{\psi\bar{\psi}}{\mu^2} - \xi). \quad (8.29)$$

Another way for calculating of kählerian effective potential consists in summarizing of contributions from supergraphs given by Fig. 4. Sum of these contributions looks like [29]

$$K^{(1)}=\int\frac{d^4k}{(2\pi)^4}\int d^4\theta_1\dots d^4\theta_{2n}\sum_{n=1}^{\infty}\frac{1}{2n}(\frac{\psi\bar{\psi}}{k^4})\frac{D^2}{4}\delta_{12}\frac{\bar{D}^2}{4}\delta_{23}\dots\frac{D^2}{4}\delta_{n-1,n}\frac{\bar{D}^2}{4}\delta_{n1}. \tag{8.30}$$

which after D-algebra transformations and summation looks like

$$K^{(1)}=\int\frac{d^{4-\epsilon}k}{(2\pi)^{4-\epsilon}}\frac{1}{2k^2}\log(1+\frac{\psi\bar{\psi}}{k^2}) \tag{8.31}$$

(here we carried out dimensional regularization by introducing parameter ϵ). Integration leads to

$$K^{(1)}=\frac{1}{32\pi^2}[\frac{\psi\bar{\psi}}{\epsilon}-\psi\bar{\psi}\log\frac{\psi\bar{\psi}}{e\mu^2}] \tag{8.32}$$

where $e=\exp(1)$. Subtraction of divergence and redefinition of μ leads to result (8.29).

Now we turn to calculation of chiral effective potential. It is not equal to zero for massless theories. Really, as it was noted by West [16] the mechanism of arising chiral corrections is the following one. If the theory describes dynamics of chiral and antichiral superfields, then quantum correction of the form

$$\int d^8z f(\Phi)(-\frac{D^2}{4\Box})g(\Phi) \tag{8.33}$$

can be rewritten as

$$\int d^6z f(\Phi)g(\Phi). \tag{8.34}$$

Here we used properties $\int d^8z=\int d^6z(-\frac{D^2}{4})$ and $\bar{D}^2D^2\Phi=16\Box\Phi$ (last identity is true for any chiral superfield Φ), and $f(\Phi)$, $g(\Phi)$ are arbitrary functions of chiral superfield Φ. However, presence of factor $\Box^{-1}$ is characteristic for massless theories, in massive theories where we have $(\Box-m^2)^{-1}$ instead of $\Box^{-1}$, and this mechanism of arising contributions to chiral effective potential is not valid. In the case of massless theory we can find matrix superpropagator exactly: first,

$$\begin{aligned}G^{\psi}_v(z_1,z_2)&=(\Box+\frac{1}{4}\psi\bar{D}^2)^{-1}\delta^8(z_1-z_2)=\frac{1}{\Box_1}\delta^8(z_1-z_2)-\\&-\frac{1}{4\Box_1}\psi(z_1)\frac{\bar{D}^2_1}{4\Box_1}\delta^8(z_1-z_2)\end{aligned} \tag{8.35}$$

(higher terms in this expansion are equal to zero because they are proportional to $\bar{D}^2\psi=0$ or $\bar{D}^4=0$). Therefore components of matrix superpropagator look like

$$\begin{aligned}G_{++}&=0; G_{+-}=G^*_{-+}=\frac{\bar{D}^2_1D^2_2}{16\Box}\delta^8(z_1-z_2)\\G_{--}&=-\frac{D^2_1}{4\Box_1}[\psi(z_1)\frac{\bar{D}^2_1D^2_2}{16\Box}\delta^8(z_1-z_2)].\end{aligned} \tag{8.36}$$

Here $*$ denotes complex conjugation. We note that background chiral superfield ψ is not constant, otherwise when we arrive at expression proportional to $D^2\psi$ we get singularity $\frac{0}{0}$ [21]. The only two-loop contribution to chiral effective potential is given by the following supergraph



$Fig.6$

External lines are chiral. We use representation in which $\Phi(z) = \Phi(x, \theta)$, and $\bar{D}_{\dot\alpha} = -\frac{\partial}{\partial\theta^{\dot\alpha}}$. Remind that in this case $\psi = \lambda\Phi$.

Contribution of the supergraph given in Fig.6 looks like

$$\begin{aligned} I &= \frac{\lambda^5}{12}\int\frac{d^4p_1d^4p_2}{(2\pi)^8}\frac{d^4kd^4l}{(2\pi)^8}\int d^4\theta_1d^4\theta_2d^4\theta_3d^4\theta_4d^4\theta_5\Phi(-p_1,\theta_3)\Phi(-p_2,\theta_4)\times \\ &\times\ \Phi(p_1+p_2,\theta_5)\frac{1}{k^2l^2(k+p_1)^2(l+p_2)^2(l+k)^2(l+k+p_1+p_2)^2}\times \\ &\times\ \delta_{13}\frac{\bar{D}_3^2}{4}\delta_{32}\frac{D_1^2\bar{D}_4^2}{16}\delta_{14}\delta_{42}\frac{D_1^2\bar{D}_5^2}{16}\delta_{15}\delta_{52}. \end{aligned} \tag{8.37}$$

After D-algebra transformation this expression can be written as

$$\begin{aligned} I &= \frac{\lambda^5}{12}\int\frac{d^4p_1d^4p_2}{(2\pi)^8}\frac{d^4kd^4l}{(2\pi)^8}\int d^2\theta\Phi(-p_1,\theta)\Phi(-p_2,\theta)\Phi(p_1+p_2,\theta)\times \\ &\times\ \frac{k^2p_1^2+l^2p_2^2+2(kl)(p_1p_2)}{k^2l^2(k+p_1)^2(l+p_2)^2(l+k)^2(l+k+p_1+p_2)^2}. \end{aligned} \tag{8.38}$$

Here we made transformation $\int d^4\theta = \int d^2\theta(-\frac{1}{4}\bar{D}^2)$ and took into account that $\bar{D}^2D^2\Phi(p,\theta) = -16p^2\Phi(p,\theta)$. Note that if $\Phi = const$ we get $\bar{D}^2D^2\Phi = 0$, hence we cannot consider Φ as constant.

As we know the effective potential is the effective lagrangian for superfields slowly varying in space-time. Let us study behaviour of the expression (8.38) in this case. The contribution (8.38) can be expressed as

$$I = \frac{\lambda^5}{12}\int d^2\theta\int\frac{d^4p_1d^4p_2}{(2\pi)^8}\Phi(-p_1,\theta)\Phi(-p_2,\theta)\Phi(p_1+p_2,\theta)S(p_1,p_2). \tag{8.39}$$

Here p_1, p_2 are external momenta. The expression $S(p_1,p_2)$ here is equal to

$$\int\frac{d^4kd^4l}{(2\pi)^8}\frac{k^2p_1^2+l^2p_2^2+2(kl)(p_1p_2)}{k^2l^2(k+p_1)^2(l+p_2)^2(l+k)^2(l+k+p_1+p_2)^2}.$$

After Fourier transform eq. (8.39) has the form

$$\begin{aligned} I &= \frac{\lambda^5}{12}\int d^2\theta \int d^4x_1 d^4x_2 d^4x_3 \int \frac{d^4p_1 d^4p_2}{(2\pi)^8}\Phi(x_1,\theta)\Phi(x_2,\theta)\times \\ &\times \Phi(x_3,\theta)\exp[i(-p_1x_1 - p_2x_2 + (p_1+p_2)x_3)]S(p_1,p_2). \end{aligned} \tag{8.40}$$

Since superfields in the case under consideration are slowly varying in space-time we can put $\Phi(x_1,\theta)\Phi(x_2,\theta)\Phi(x_3,\theta) \simeq \Phi^3(x_1,\theta)$.. As a result one gets

$$\begin{aligned} I &= \frac{\lambda^5}{12}\int d^2\theta \int d^4x_1 d^4x_2 d^4x_3 \int \frac{d^4p_1 d^4p_2}{(2\pi)^8}\Phi^3(x_1,\theta)\times \\ &\times \exp[i(-p_1x_1 - p_2x_2 + (p_1+p_2)x_3)]S(p_1,p_2). \end{aligned} \tag{8.41}$$

Integration over $d^4x_2 d^4x_3$ leads to delta-functions $\delta(p_2)\delta(p_1+p_2)$. Hence the eq. (8.40) takes the form

$$I = \frac{\lambda^5}{12}\int d^2\theta \int d^4x_1 \Phi^3(x_1,\theta) S(p_1,p_2)|_{p_1,p_2=0}. \tag{8.42}$$

Therefore final result for two-loop correction to chiral (frequently called holomorphic) effective potential looks like

$$W^{(2)} = \frac{6}{(16\pi^2)^2}\zeta(3)\Phi^3(z). \tag{8.43}$$

Here we took into account that

$$\int \frac{d^4k d^4l}{(2\pi)^8}\frac{k^2p_1^2 + l^2p_2^2 + 2(k_1k_2)(p_1p_2)}{k^2l^2(k+p_1)^2(l+p_2)^2(l+k)^2(l+k+p_1+p_2)^2}|_{p_1=p_2=0} = \frac{6}{(4\pi)^4}\zeta(3).$$

We see that the correction (8.43) is finite and does not require renormalization.

Chiral contributions to effective action arise also in other theories describing dynamics of chiral superfields. F.e. in general chiral superfield theory the leading chiral contribution is also chiral effective potential (see next section), in dilaton supergravity leading chiral contribution is of second order in space-time derivatives of chiral superfield (see section 10), and these corrections are finite. The situation in $N = 1$ super-Yang-Mills theory, however, possesses some peculiarities. Really, in this model both finite (Fig. 7a) and divergent (Fig.7b) two-loop chiral contributions are possible.



Fig.7a



Fig.7b

Finite contributions (there are more than 10 supergraphs of the form similar to Fig. 7a [30]) all give contributions to effective potential proportional to $\frac{\zeta(3)}{(4\pi)^4}\Phi^3$ (cf. (8.43)), i.e. finite chiral contributions are analogous to the case of Wess-Zumino model. As for divergent contribution given by Fig. 4b it is, after D-algebra transformation, equal to

$$\int d^2\theta \int \frac{d^4p_1 d^4p_2}{(2\pi)^8}(p_1+p_2)^2 \int \frac{d^4k d^4l}{(2\pi)^8}\frac{1}{k^2(k+p_1)^2(k+p_2)^2} \times$$
$$\times \quad \frac{1}{l^2(l+p_1+p_2)^2}\Phi(-p_1,\theta)\Phi(-p_2,\theta)\Phi(p_1+p_2,\theta). \tag{8.44}$$

To obtain the low-energy leading contribution we must consider the limit at $p_1, p_2 \to 0$. It is known [20] that

$$\lim_{p_1,p_2\to 0}(p_1+p_2)^2\int\frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p_1)^2(k+p_2)^2} = \frac{1}{16\pi^2}\int_0^1 d\alpha\frac{\log[\alpha(1-\alpha)]}{1-\alpha(1-\alpha)} = \frac{C_0}{16\pi^2},$$

where C_0 is a some constant. The integral over l is divergent, and after dimensional regularization it is equal to

$$\int\frac{d^{4-\epsilon}l}{(2\pi)^{4-\epsilon}}\frac{1}{l^2(l+p_1+p_2)^2} = \frac{1}{16\pi^2}(\frac{2}{\epsilon}+\log\frac{(p_1+p_2)^2}{\mu^2}).$$

After cancellation of divergence with help of the appropriate one-loop counterterm and transforming to coordinate representation we see that expression (8.44) for slowly varying in space-time superfields takes the form

$$\int d^6z\frac{1}{(16\pi^2)^2}\Phi^2(z)\log(-\frac{\Box}{\mu^2})\Phi(z). \tag{8.45}$$

Thus, the leading chiral correction in $N=1$ super-Yang-Mills theory with chiral matter is nonlocal one (detailed discussion is given in [30]). We note that nonlocal chiral corrections in this theory arise also in the pure gauge sector [31]. Hence presence of quantum contributions to chiral effective Lagrangian is quite characteristic for theories including chiral superfields.

9 General chiral superfield model

From viewpoint of superstring theory low-energy models of elementary particles are effective theories in which integration over massive string modes is carried out, 10-dimensional background manifold has the form $M^4\times K$ where M^4 is four-dimensional Minkovski space, and K is some six-dimensional compact manifold. Then reduction to M^4 is carried out. As a result we arrive at theory in M^4 with action [1]

$$S[\Phi,\bar\Phi] = \int d^8zK(\Phi^i,\bar\Phi^i) + (\int d^6zW(\Phi^i)+h.c.). \tag{9.1}$$

We can use matrix denotions via introduction of column vector $\vec\Phi = \{\Phi^i\}$, after which the consideration in the case of several chiral superfield is analogous to the case of one chiral superfield (some extensions of this model involving the gauge superfields are given in [32]). We can consider this theory for arbitrary functions K and W. Note that there is no higher derivatives in the classical action. Therefore the theory with the action (9.1) is the most general theory without higher derivatives describing dynamics of chiral superfield. There are a lot of phenomenological applications of this model in string theory (see [32] and references therein). In general case this theory is nonrenormalizable; however, it is an effective theory aimed for studying of the low-energy domain. Therefore all integrals over momenta are effectively cut by condition $p \ll M_{String}$ where p is momentum, and $M_{String} = 10^{17} GeV \sim 10^{-2} M_{Pl}$ is a characteristic string mass.

The effective action in the theory, as well as that one in the Wess-Zumino model, can be presented as a series in supercovariant derivatives $D_A = (\partial_a, D_\alpha, \bar{D}_{\dot\alpha})$ in the form

$$\begin{aligned}\Gamma[\bar\Phi, \Phi] &= \int d^8 z \mathcal{L}_{eff}(\Phi, D_A\Phi, D_A D_B\Phi; \bar\Phi, D_A\bar\Phi, D_A D_B\bar\Phi) + \\ &+ \ (\int d^6 z \mathcal{L}^{(c)}_{eff}(\Phi) + h.c.) + \ldots . \end{aligned} \tag{9.2}$$

Here again, like in the Wess-Zumino model, $\mathcal{L}_{eff}$ is called general effective lagrangian, $\mathcal{L}^{(c)}_{eff}$ is called chiral effective lagrangian. Both these lagrangians are the series in supercovariant derivatives of superfields and can be written in the form of loop expansion

$$\begin{aligned}\mathcal{L}_{eff} &= K_{eff}(\bar\Phi, \Phi) + \ldots = K(\bar\Phi, \Phi) + \sum_{n=1}^{\infty} K^{(n)}_{eff}(\bar\Phi, \Phi); \\ \mathcal{L}^{(c)}_{eff} &= W_{eff}(\Phi) + \ldots = W(\Phi) + \sum_{n=1}^{\infty} W^{(n)}_{eff}(\Phi) + \ldots . \end{aligned} \tag{9.3}$$

Here dots mean terms depending on covariant derivatives of superfields $\Phi, \bar\Phi$. As earlier, the $K_{eff}(\bar\Phi, \Phi)$ is called kahlerian effective potential, $W_{eff}(\Phi)$ is called chiral (or holomorphic) effective potential, $K^{(n)}_{eff}$ is a n-th correction to kahlerian potential and $W^{(n)}_{eff}$ is a n-th correcton to chiral (holomorphic) potential W.

The one-loop contribution to the effective action is totally determined by the quadratic part of expansion of $\frac{1}{\hbar} S[\bar\Phi + \sqrt{\hbar}\bar\phi, \Phi + \sqrt{\hbar}\phi]$ in quantum fields $\phi, \bar\phi$ which looks like

$$S_2 = \frac{1}{2} \int d^8 z \begin{pmatrix} \phi & \bar\phi \end{pmatrix} \begin{pmatrix} K_{\Phi\Phi} & K_{\Phi\bar\Phi} \\ K_{\Phi\bar\Phi} & K_{\bar\Phi\bar\Phi} \end{pmatrix} \begin{pmatrix} \phi \\ \bar\phi \end{pmatrix} + [\int d^6 z \frac{1}{2} W'' \phi^2 + h.c.] \tag{9.4}$$

and defines the propagators and the the higher terms of expansion define the vertices. Here $K_{\Phi\bar\Phi} = \frac{\partial^2 K(\bar\Phi, \Phi)}{\partial\Phi\partial\bar\Phi}$, $K_{\Phi\Phi} = \frac{\partial^2 K(\bar\Phi,\Phi)}{\partial\Phi^2}$ etc, $W'' = \frac{d^2 W}{d\Phi^2}$.

The corresponding matrix superpropagator has the form

$$G(z_1, z_2) = \begin{pmatrix} G_{++}(z_1, z_2) & G_{+-}(z_1, z_2) \\ G_{-+}(z_1, z_2) & G_{--}(z_1, z_2) \end{pmatrix} \tag{9.5}$$

where $+$ denotes chirality with respect to corresponding argument, and $-$ correspondingly – antichirality. This propagator satisfies the equation

$$\begin{pmatrix} W'' - \frac{1}{4}(\bar{D}^2 K_{\Phi\Phi}) & -\frac{1}{4}\vec{\bar{D}}^2 K_{\Phi\bar\Phi} \\ -\frac{1}{4}\vec{D}^2 K_{\Phi\bar\Phi} & \bar{W}'' - \frac{1}{4}(D^2 K_{\bar\Phi\bar\Phi}) \end{pmatrix} \begin{pmatrix} G_{++}(z_1, z_2) & G_{+-}(z_1, z_2) \\ G_{-+}(z_1, z_2) & G_{--}(z_1, z_2) \end{pmatrix} =$$

$$= \; -\begin{pmatrix} \delta_+ & 0 \\ 0 & \delta_- \end{pmatrix}. \tag{9.6}$$

To consider kählerian effective potential we must omit all derivatives of superfields $\Phi, \bar{\Phi}$, and the equation takes the form

$$\begin{pmatrix} W'' & -\frac{1}{4}K_{\Phi\bar{\Phi}}\bar{D}^2 \\ -\frac{1}{4}K_{\Phi\bar{\Phi}}D^2 & \bar{W}'' \end{pmatrix}\begin{pmatrix} G_{++}(z_1,z_2) & G_{+-}(z_1,z_2) \\ G_{-+}(z_1,z_2) & G_{--}(z_1,z_2) \end{pmatrix} = -\begin{pmatrix} \delta_+ & 0 \\ 0 & \delta_- \end{pmatrix}. \tag{9.7}$$

The solution of this equation looks like

$$G = \frac{1}{K^2_{\Phi\bar{\Phi}}\Box - W''\bar{W}''}\begin{pmatrix} W'' & \frac{1}{4}K_{\Phi\bar{\Phi}}\bar{D}^2 \\ \frac{1}{4}K_{\Phi\bar{\Phi}}D^2 & \bar{W}'' \end{pmatrix}\begin{pmatrix} \delta_+ & 0 \\ 0 & \delta_- \end{pmatrix}. \tag{9.8}$$

Now we turn to studying of quantum contributions to kählerian effective potential depending only on superfields Φ, $\bar{\Phi}$ but not on their derivatives.

The one-loop diagrams contributing to kahlerian effective potential are







. . .

Double external lines correspond to alternating W'' and $\bar{W}''$. Internal lines are $<\phi\bar{\phi}>$-propagators of the form

$$G_0 \equiv <\phi\bar{\phi}> = -\frac{\bar{D}^2 D^2}{16K_{\Phi\bar{\Phi}}\Box}.$$

Supergraph of such structure with $2n$ legs represents itself as a ring containing n links of the following form



The total contribution of all these diagrams after D-algebra transformations, summation, integration over momenta and subtraction of divergences is equal to

$$K^{(1)} = -\int d^4\theta \mathrm{tr}\frac{1}{32\pi^2}W''\frac{1}{K^2_{\Phi\bar{\Phi}}}\bar{W}''\ln\Big(W''\frac{1}{\mu^2 K^2_{\Phi\bar{\Phi}}}\bar{W}''\Big). \tag{9.9}$$

This form is more convenient for analysis of many-field model than that one given in [12, 33], and tr denotes trace of product of the given matrices. It is easy to show that the present result corresponds to the known expression for the Wess-Zumino model where $W'' = \frac{1}{2}\lambda\Phi$.

Let us consider the chiral (holomorphic) effective potential $W_{eff}(\Phi)$. The mechanism of its arising is just the same than in Wess-Zumino model. We note again that the chiral contributions to effective action can be generated by supergraphs containing massless propagators only. To find chiral corrections to effective action we put $\bar{\Phi} = 0$ in eq. (9.4). **Therefore here and further all derivatives of K, W and $\bar{W}$ will be taken at $\bar{\Phi} = 0$.** Under this condition the action of quantum superfields $\phi, \bar{\phi}$ in external superfield Φ looks like

$$S[\bar{\phi}, \phi, \Phi] = \frac{1}{2}\int d^8z \begin{pmatrix} \phi & \bar{\phi} \end{pmatrix} \begin{pmatrix} K_{\Phi\Phi} & K_{\Phi\bar{\Phi}} \\ K_{\Phi\bar{\Phi}} & K_{\bar{\Phi}\bar{\Phi}} \end{pmatrix} \begin{pmatrix} \phi \\ \bar{\phi} \end{pmatrix} + \int d^6z \frac{1}{2} W^{''} \phi^2 + \ldots . \qquad (9.10)$$

The dots here denote the terms of third, fourth and higher orders in quantum superfields. We call the theory massless if $W^{''}|_{\Phi=0} = 0$. Further we consider only massless theory.

To calculate the corrections to $W(\Phi)$ we use supergraph technique (see f.e. [10]). For this purpose one splits the action (9.10) into sum of free part and vertices of interaction. As a free part we take the action $S_0 = \int d^8z\phi\bar{\phi}$. The corresponding superpropagator is $G(z_1, z_2) = -\frac{D_1^2\bar{D}_2^2}{16\Box}\delta^8(z_1 - z_2)$. And the term $S[\bar{\phi}, \phi, \Phi] - S_0$ will be treated as vertices where $S[\bar{\phi}, \phi, \Phi]$ is given by eq. (9.10). Our purpose is to find the first leading contribution to $W_{eff}(\Phi)$. As we will show, chiral loop contributions begin with two loops. Therefore we keep in eq. (9.10) only the terms of second, third and fourth orders in quantum fields.

Non-trivial corrections to chiral potential can arise only if $2L + 1 - n_{W''} - n_{V_c} = 0$ where L is a number of loops, $n_{W''}$ is a number of vertices proportional to $W^{''}$, n_{V_c} is that one of vertices of third and higher orders in quantum superfields, otherwise corresponding contribution will either vanish or lead to singularity in the infrared linit. In one-loop approximation this equation leads to $n_{W''} + n_{V_c} = 3$. However, all supergraphs satisfying this condition have zero contribution. Therefore first correction to chiral effective potential is two-loop one. In two-loop approximation this equation has the form $n_{W''} + n_{V_c} = 5$. Since the number of purely chiral (antichiral) vertices independent of $W^{''}$ in two-loop supergraphs can be equal to 0, 1 or 2, the number of external vertices $W^{''}$ takes values from 3 to 5.

We note that non-trivial contribution to chiral (holomorphic) effective potential from any diagram can arise only if the number of D^2-factors is more by one than the number of $\bar{D}^2$-factors (see details in [34]). The only Green function in the theory is the propagator $< \phi\bar{\phi} >$. Therefore total number of quantum chiral superfields ϕ corresponding to all vertices must be equal to that one of antichiral ones $\bar{\phi}$. As a result we find that the only two-loop supergraph contributing to chiral effective potential looks like



*Fig.*8

Here internal lines are propagators $<\phi\bar{\phi}>$ depending on background chiral superfields which have the form

$$<\phi\bar{\phi}>= -\bar{D}_1^2 D_2^2 \frac{\delta^8(z_1-z_2)}{16K_{\Phi\bar{\Phi}}(z_1)\Box}. \tag{9.11}$$

We note that the superfield $K_{\Phi\bar{\Phi}}$ is not constant here. Double external lines are W''.

After D-algebra transformations and loop integrations we find that two-loop contribution to holomorphic effective potential in this model looks like

$$W^{(2)} = \frac{6}{(16\pi^2)^2}\zeta(3)\bar{W}'''^2\Big\{\frac{W''(z)}{K^2_{\Phi\bar{\Phi}}(z)}\Big\}^3. \tag{9.12}$$

One reminds that $\bar{W}''' = \bar{W}'''(\bar{\Phi})|_{\bar{\Phi}=0}$ and $K_{\Phi\bar{\Phi}}(z) = \frac{\partial^2 K(\bar{\Phi},\Phi)}{\partial\Phi\partial\bar{\Phi}}|_{\bar{\Phi}=0}$ here. We see that the correction (9.12) is finite and does not require renormalization in any case despite the theory is non-renormalizable in general case.

We note that the calculation of two-loop kählerian effective potential can be carried out with help of matrix superpropagator (9.8). The results are given in [12, 33].

Now let us consider some phenomenological applications of the theory characterized by the action (9.1). Let us suppose that the column vector $\vec{\Phi}$ describes two superfield: light (massless) ϕ and heavy Φ, $\vec{\Phi} = \begin{pmatrix} \phi \\ \Phi \end{pmatrix}$. We calculate for this case one-loop effective action and eliminate heavy superfields with use of effective equations of motion. As a result we arrive at the effective action of light superfields. There is a decoupling theorem [18, 35] according to which this effective action after redefining of parameters (fields, masses, coupling) can be expressed in the form of a sum of effective action of the theory obtained from initial one by putting heavy fields to zero and terms proportional to different powers of $\frac{1}{M}$ where M is mass of heavy superfield (which in the case under consideration is put, by fenomenological reasons, to be equal to M_{String} [32, 33]).

We study such a theory in one-loop approximation. The low-energy leading one-loop contribution to effective action is given by (9.9). The matrices $K_{\Phi\bar{\Phi}} = \frac{\partial^2 K}{\partial\Phi^i\partial\bar{\Phi}^j}$ and $W'' = \frac{\partial^2 W}{\partial\Phi^i\partial\Phi^j}$ can be diagonalized simultaneously, and the trace is a sum of proper values.

We consider, as an example, the minimal theory [36] with

$$K \;=\; \phi\bar{\phi} + \Phi\bar{\Phi};\; W = \frac{M}{2}\Phi^2 + \frac{\lambda}{2}\Phi\phi^2 + \frac{g}{3!}\phi^3. \tag{9.13}$$

After calculations described in [36] and analogous ones to those of Wess-Zumino model we get

$$K^{(1)} \;=\; -\frac{1}{32\pi^2}(R_1^2\log\frac{R_1^2}{\mu^2} + R_2^2\log\frac{R_2^2}{\mu^2}) \tag{9.14}$$

where

$$\begin{aligned} R_{1,2} \;&=\; |\lambda\Phi + g\phi|^2 + 2\lambda^2|\phi|^2 + M^2 \pm \\ &\pm\; \sqrt{(|\lambda\Phi + g\phi|^2 - M^2)^2 + 4|\lambda^2\Phi\bar{\phi} + \lambda M\phi + \lambda g|\phi|^2|^2}. \end{aligned} \tag{9.15}$$

The low-energy effective action is given by

$$\Gamma^{(1)} \;=\; \int d^8z(\phi\bar{\phi} + \Phi\bar{\Phi} + \hbar K^{(1)}) + [\int d^6z(\frac{M}{2}\Phi^2 + \frac{\lambda}{2}\Phi\phi^2 + \frac{g}{3!}\phi^3) + h.c.]. \tag{9.16}$$

The effective equations of motion for heavy superfield Φ looks like

$$-\frac{1}{4}\bar{D}^2(\bar{\Phi} + \hbar\frac{\partial K^{(1)}}{\partial\Phi}) + M\Phi + \frac{\lambda}{2}\phi^2 = 0. \tag{9.17}$$

We can solve this equation by iterative method, i.e. we suppose that

$$\Phi = \Phi_0 + \Phi_1 + \ldots + \Phi_n + \ldots, \tag{9.18}$$

with Φ_k is k-th approximation. Since mass M is very large we suppose that $|D^2\Phi| \ll M\Phi$. Zero approximation is obtained from condition $M\Phi_0 + \frac{\lambda}{2}\phi^2 = 0$, i.e. $\Phi_0 = -\frac{\lambda\phi^2}{2M}$. And k-th approximation is proportional to M^{-k}. Substituting the expansion of Φ (9.18) into (9.17) we get the following recurrent relation for Φ_{n+1}:

$$\Phi_{n+1} = \frac{\bar{D}^2}{4M}[\bar{\Phi}_n + \hbar\frac{\partial K^{(1)}}{\partial\Phi}|_{\Phi=\Phi_0+\ldots+\Phi_n} - \hbar\frac{\partial K^{(1)}}{\partial\Phi}|_{\Phi=\Phi_0+\ldots+\Phi_{n-1}}]. \tag{9.19}$$

As a result, we get the following solution for Φ in leading order

$$\Phi = -\frac{\lambda\phi^2}{2M} - \frac{\hbar\bar{D}^2}{64\pi^2 M}[\lambda g\bar{\phi}(1 + \log\frac{2g^2|\phi|^2}{\mu^2})] + O(\frac{1}{M^2}). \tag{9.20}$$

This solution can be substituted into the low-energy effective action (9.16). The effective action of light superfields is defined as

$$S_{eff}[\phi,\bar{\phi}] = \Gamma^{(1)}[\phi,\bar{\phi};\Phi(\phi,\bar{\phi}),\bar{\Phi}(\phi,\bar{\phi})] \tag{9.21}$$

where $\Phi(\phi,\bar{\phi})$ and the same notation for $\bar{\Phi}$ mean that heavy superfields are expressed in terms of light ones via effective equations of motion. In the case under consideration, effective action of light superfields looks like

$$S_{eff} = S^{(0)}_{eff} + S^{(1)}_{eff} + O(\frac{1}{M^2}) \tag{9.22}$$

where $S^{(0)}_{eff}$ and $S^{(1)}_{eff}$ are of zeroth and first orders in M^{-1} respectively. These expressions look like

$$\begin{aligned}
S^0_{eff} &= \int d^8z\Big[\phi\bar{\phi} - \frac{\hbar}{32\pi^2}[4\lambda^2|\phi|^2(1+\log\frac{2M^2}{\mu^2}) + 2g^2|\phi|^2\log\frac{2g^2|\phi|^2}{\mu^2}] + \\
&+ (\int d^6z\frac{g}{3!}\phi^3 + h.c.)\Big] \\
S^{(1)}_{eff} &= -\frac{\hbar}{32\pi^2M}\int d^8z\Big[-2\lambda^2g(\phi+\bar{\phi})|\phi|^2\log\frac{g^2|\phi|^2}{M^2} + 2\lambda^2g[\bar{\phi}\phi^2 + h.c.]\Big] + \\
&+ \frac{1}{M}\Big[\int d^6z\Big(\frac{1}{8}\lambda^2\phi^4 - \lambda^2g^2(\frac{\hbar\bar{D}^2}{64\pi^2}(\bar{\phi}(1+\log\frac{2g^2|\phi|^2}{\mu^2}))^2\Big) + h.c.\Big].
\end{aligned} \tag{9.23}$$

We see that this effective action contains term of the form

$$-\frac{\hbar}{32\pi^2}4\lambda^2|\phi|^2(1+\log\frac{2M^2}{\mu^2})$$

which increases with growth of M. If we put the renormalization condition $\log\frac{2M^2}{\mu^2} = -1$ this term vanishes. The effective action, however, in this case takes the form

$$\begin{aligned}
S^0_{eff} &= \int d^8z\Big[\phi\bar{\phi} - \frac{\hbar}{32\pi^2}2g^2|\phi|^2\log\frac{2g^2|\phi|^2}{eM^2} + \\
&+ (\int d^6z\frac{g}{3!}\phi^3 + h.c.)\Big].
\end{aligned} \tag{9.24}$$

This expression contains the term $2g^2|\phi|^2\log\frac{2g^2|\phi|^2}{eM^2}$ which also increases as M grows. Therefore we see that modifications of light superfield effective action caused by presence of heavy superfields are significant [36].

10 Dilaton supergravity as an example of superfield theory with higher derivatives

The starting point of our consideration is the supertrace anomaly of matter superfield in curved superspace. The action generating this anomaly is obtained in [37]. In conformally flat superspace (in which the vector supergravity prepotential H^m is equal to zero) this action looks like

$$\begin{aligned}
\Gamma_A &= \frac{1}{16\pi^2}\int d^8z(8c\partial_a\bar{\sigma}\partial^a\sigma + \frac{1}{32}(32c-b)\bar{D}^{\dot{\alpha}}\bar{\sigma}D^{\alpha}\sigma(\partial_{\alpha\dot{\alpha}}(\sigma+\bar{\sigma}) - \\
&- \frac{1}{2}\bar{D}_{\dot{\alpha}}\bar{\sigma}D_{\alpha}\sigma)).
\end{aligned} \tag{10.1}$$

Here we have used the "flat" supercovariant derivatives D_α, $\bar{D}_{\dot\alpha}$, $\partial_{\alpha\dot\alpha}$; and $\sigma = \ln\Phi$, $\bar\sigma = \ln\bar\Phi$.

Let us also consider the superfield action of N=1 supergravity, in conformally flat superspace we obtain

$$S_{SG} = -\frac{m^2}{2}\int d^8z\Phi\bar\Phi + [\Lambda\int d^6z\Phi^3 + h.c.], \tag{10.2}$$

where $m^2 = \frac{6}{\kappa^2}$, Λ is the cosmological constant and κ is the gravitational constant. We will investigate the theory action of which is a sum of the actions (10.1) and (10.2) Denoting

$$\frac{Q^2}{2} = 8c,\ \xi_1 = \frac{1}{32(4\pi)^2}(32c - b),\ \xi_2 = -\frac{1}{64(4\pi)^2}(32c - b),$$

we get a superfield theory in flat superspace with the action of the form

$$\begin{aligned} S &= \int d^8z(-\frac{Q^2}{2(4\pi)^2}\bar\sigma\Box\sigma + \bar{D}^{\dot\alpha}\bar\sigma D^\alpha\sigma \times \\ &\times (\xi_1\partial_{\alpha\dot\alpha}(\sigma+\bar\sigma) + \xi_2\bar{D}_{\dot\alpha}\bar\sigma D_\alpha\sigma) - \frac{m^2}{2}e^{\sigma+\bar\sigma}) + (\Lambda\int d^6z e^{3\sigma} + h.c.). \end{aligned} \tag{10.3}$$

The $Q^2, \xi_1, \xi_2, m^2, \Lambda$ will be considered as the arbitrary and independent parameters of the model. We will call the model with action (10.3) the four-dimensional dilaton supergravity model.

In order to calculate the counterterms and to find the divergences we should study the structure of supergraphs of the theory. The strucure of supergraphs is defined by a form of propagators and vertices.

The theory under consideration is characterized by a matrix propagator

$$G(z,z') = \begin{pmatrix} G_{++}(z,z') & G_{+-}(z,z') \\ G_{-+}(z,z') & G_{--}(z,z') \end{pmatrix}, \tag{10.4}$$

satisfying the equation

$$\begin{pmatrix} 9\Lambda & (\frac{Q^2}{(4\pi)^2}\Box + m^2)\frac{\bar{D}^2}{4} \\ (\frac{Q^2}{(4\pi)^2}\Box + m^2)\frac{D^2}{4} & 9\bar\Lambda \end{pmatrix}\begin{pmatrix} G_{++} & G_{+-} \\ G_{-+} & G_{--} \end{pmatrix} = \begin{pmatrix} \delta_+ & 0 \\ 0 & \delta_- \end{pmatrix}, \tag{10.5}$$

where $\delta_+ = -\frac{1}{4}\bar{D}^2\delta^8(z_1 - z_2)$, $\delta_- = -\frac{1}{4}D^2\delta^8(z_1 - z_2)$. The solution of this equation is written in the form

$$G = \frac{1}{81\Lambda\bar\Lambda - \Box(\frac{Q^2}{16\pi^2}\Box + m^2)^2}\begin{bmatrix} 9\bar\Lambda & -(\frac{Q^2}{16\pi^2}\Box + m^2)\frac{\bar{D}^2}{4} \\ -(\frac{Q^2}{16\pi^2}\Box + m^2)\frac{D^2}{4} & 9\Lambda \end{bmatrix}. \tag{10.6}$$

This propagator acts on columns $\begin{pmatrix} \phi \\ \bar\phi \end{pmatrix}$, where ϕ is a chiral superfield and $\bar\phi$ is an antichiral superfield.

The propagator in momentum representation looks like follows

$$\begin{aligned} G_{++}(k) &= \bar{G}_1(k)\frac{-\bar{D}^2}{4}\delta_{12}\, G_{+-}(k) = G_2(k)\frac{\bar{D}^2 D^2}{16}\delta_{12} \\ G_{-+}(k) &= G_2(k)\frac{D^2\bar{D}^2}{16}\delta_{12}\, G_{--}(k) = G_1(k)\frac{-D^2}{4}\delta_{12}, \end{aligned} \tag{10.7}$$

where

$$\begin{aligned} G_1(k) &= \frac{9\Lambda}{k^2(\frac{Q^2}{16\pi^2}k^2 - m^2)^2 + 81\Lambda\bar{\Lambda}}\, \bar{G}_1(k) = \frac{9\bar{\Lambda}}{k^2(\frac{Q^2}{16\pi^2}k^2 - m^2)^2 + 81\Lambda\bar{\Lambda}} \\ G_2(k) &= \frac{-\frac{Q^2}{16\pi^2}k^2 + m^2}{k^2(\frac{Q^2}{16\pi^2}k^2 - m^2)^2 + 81\Lambda\bar{\Lambda}} \end{aligned}$$

A structure of vertices is taken from action (10.3). There are four vertices:

$$\begin{aligned} V_1 &= \xi_1 \bar{D}^{\dot{\alpha}}\bar{\sigma} D^{\alpha}\sigma\partial_{\alpha\dot{\alpha}}(\sigma+\bar{\sigma}); \\ V_2 &= \xi_2 \bar{D}_{\dot{\alpha}}\bar{\sigma}\bar{D}^{\dot{\alpha}}\bar{\sigma} D^{\alpha}\sigma D_{\alpha}\sigma; \\ V_3 &= -\frac{m^2}{2}(e^{\sigma+\bar{\sigma}} - 1 - (\sigma+\bar{\sigma}) - \frac{1}{2}(\sigma+\bar{\sigma})^2); \\ V_4 &= \Lambda(-\frac{D^2}{4\Box})(e^{3\sigma} - 1 - 3\sigma - \frac{9}{2}\sigma^2) + \bar{\Lambda}(-\frac{\bar{D}^2}{4\Box})(e^{3\bar{\sigma}} - 1 - 3\bar{\sigma} - \frac{9}{2}\bar{\sigma}^2). \end{aligned} \tag{10.8}$$

The factors $(-\frac{D^2}{4\Box})$ and $(-\frac{\bar{D}^2}{4\Box})$ arise in V_4 since the all vertices correspond to the action written as the integrals over whole superspace.

The eqs. (10.7–10.8) are the basis of supergraph technique allowing to develop a perturbative treatment for the model (10.3).

Let us consider the superficial degree of divergence (SDD) from this model. Since space-time derivatives give contribution to SDD equal to 1, and spinor ones – to 1/2 we see that any V_1, V_2-type vertex contributes to SDD with 2, and all such vertices – with $V_{1,2}$ (here $V_{1,2}$ is a number of such vertices). Any loop as usual contributes with 2 (4, because any loop includes integration over d^4k, and -2, since contraction of any loop into a point requires four D-factors and reduces possible contribution to SDD by 2), hence all loops – with 2L. Contribution of all propagators G_{++}, G_{--} is equal to $-5P_1$ where P_1 is a number of such propagators, and contribution of all propagators G_{+-}, G_{-+} – to $-2P_2$. The V_3-vertices do not contribute at all, and V_4-type totalize a contribution of $-V_4$ [13]. We note also that any D-factor acting to the external line instead of the internal one decreases the SDD by $\frac{1}{2}$, and any space-time derivative acting on the external line – by 1. Therefore total SDD is equal to

$$\omega = 2L + 2V_{1,2} - 5P_1 - 2P_2 - V_4 - \frac{1}{2}N_D - N_{\partial}.$$

Since $V_{1,2} + V_3 + V_4 = V$ and $L + V - P = 1$ we get

$$\omega = 2 - 3P_1 - 2V_3 - 3V_4 - \frac{1}{2}N_D - N_{\partial}. \tag{10.9}$$

This equation allows one to make some conclusions. First of all, divergent diagrams cannot contain V_4-vertices (hence all chiral contributions to effective action are finite). Second, divergent diagrams cannot contain $<\sigma\sigma>, <\bar{\sigma}\bar{\sigma}>$-propagators and can contain no more than one vertex proportional to m^2. Then, formally they can contain arbitrary number of $V_{1,2}$-vertices, hence there is an infinite number of divergent structures (f.e. divergent corrections of the form $\xi_1\xi_2^{n+1}\sigma^n\bar{\sigma}^n\bar{D}^{\dot{\alpha}}\bar{\sigma}D^{\alpha}\sigma\partial_{\alpha\dot{\alpha}}(\sigma+\bar{\sigma})$ can arise for any n). Hence the theory is non-renormalizable (in the one-loop order some of derivatives asociated to the V_1, V_2-vertices are enforced to act to the external lines which decreases the degree of divergence, in part, the graphs depicted at Figs. 9, 10 are only logarithmically divergent, but it is not valid for the higher loop orders). However, if we put $\xi_1 = \xi_2 = 0$ the theory is super-renormalizable.

Consider one-loop counterterms leading to renormalization of ξ_1, ξ_2.



Fig.9



Fig.10

The supergraphs associated with Fig.9 and Fig.10 lead to the following contributions respectively

$$\begin{aligned}
{S_1}^{(1)} &= 72\mu^{-\epsilon}\xi_1\xi_2\int d^4\theta_1 d^4\theta_2\frac{d^dp_1d^dp_2}{(2\pi)^{2d}}\times\\
&\times\ \bar{D}^{\dot{\alpha}}\bar{\sigma}(-p_1,\theta_1)D^{\alpha}\sigma(-p_2,\theta_1)\partial_{\beta\dot{\beta}}(\sigma(p,\theta_2)+\bar{\sigma}(p,\theta_2))\times\\
&\times\ \int\frac{d^dk}{(2\pi)^d}D_{\alpha}\bar{D}^{\dot{\beta}}G_{+-}(k)\bar{D}_{\dot{\alpha}}D^{\beta}G_{-+}(k+p); \qquad (10.10)\\
{S_2}^{(1)} &= 72\mu^{-\epsilon}\xi_2^2\int d^4\theta_1 d^4\theta_2\frac{d^dp_1d^dp_2d^dp_3}{(2\pi)^{3d}}\times\\
&\times\ \bar{D}^{\dot{\alpha}}\bar{\sigma}(-p_1,\theta_1)D^{\alpha}\sigma(-p_2,\theta_1)\bar{D}_{\dot{\beta}}\bar{\sigma}(p_3,\theta_2)D_{\beta}\sigma(p-p_3,\theta_2)\times\\
&\times\ \int\frac{d^dk}{(2\pi)^d}D_{\alpha}\bar{D}^{\dot{\beta}}G_{+-}(k)\bar{D}_{\dot{\alpha}}D^{\beta}G_{-+}(k+p).
\end{aligned}$$

Here ${S_1}^{(1)}$ and ${S_2}^{(1)}$ are one-loop divergent corrections to vertices V_1 and V_2 correspondingly, p_1, p_2 and p_3 are external momenta, $p = p_1+p_2$. μ is a standard arbitrary parameter of mass dimension introduced in dimensional regularization and $\epsilon = 4-d$. After calculations described in [13] we get one-loop quantum corrections from this supergraphs in the form

$$S_1^{(1)} = -576\mu^{-\epsilon}\frac{\xi_1\xi_2(16\pi^2)^2}{Q^4}\int d^8z\bar{D}^{\dot{\alpha}}\bar{\sigma}D^{\alpha}\sigma\partial_{\alpha\dot{\alpha}}(\sigma+\bar{\sigma})(\frac{2}{16\pi^2\epsilon}+fin)\equiv \qquad (10.11)$$

$$
\begin{aligned}
&\equiv S^{(1)}_{1_{div}} + S^{(1)}_{1_{fin}};\\
S_2^{(1)} &= -576\mu^{-\epsilon}\frac{\xi_2^2(16\pi^2)^2}{Q^4}\int d^8z\bar{D}^{\dot{\alpha}}\bar{\sigma}D^{\alpha}\sigma\bar{D}_{\dot{\alpha}}\bar{\sigma}D_{\alpha}\sigma(\frac{2}{16\pi^2\epsilon}+fin)\equiv\\
&\equiv S^{(1)}_{2_{div}} + S^{(1)}_{2_{fin}}.
\end{aligned}
$$

In order to renormalize the theory we introduce the one-loop counterterms $-S_1^{(1)}{}_{div}$, $-S_2^{(1)}{}_{div}$. It corresponds to the following renormalization transformation

$$
\begin{aligned}
Q^2_{(0)} &= \mu^{-\epsilon}Z_Q Q^2;\\
\xi_{1(0)} &= \mu^{-\epsilon}Z_1\xi_1;\\
\xi_{2(0)} &= \mu^{-\epsilon}Z_2\xi_2
\end{aligned}
\tag{10.12}
$$

where $Q^2_{(0)}, \xi_{1(0)}, \xi_{2(0)}$ are the bare parameters and Q^2, ξ_1, ξ_2 are the renormalized ones. As a result one obtains

$$
Z_1 = Z_2 = (1+\frac{72\xi_2(16\pi)^2}{Q^4\epsilon}).
\tag{10.13}
$$

We see that in one-loop approximation there is the same independent renormalization constant both for ξ_1 and ξ_2. It means in particular that if we put $\xi_2^{(0)} = c\xi_1^{(0)}$, where c is a constant, then the renormalized parameters ξ_1 and ξ_2 will satisfy the same relation $\xi_2 = c\xi_1$. One-loop renormalization does not destroy the relationship between the parameters in lower order.

Next step is a calculation of Z_Q. Let us consider the supergraph given on Fig.11 (note that we considered such a supergraph in Section 3)



Fig.11

The corresponding contribution looks like this

$$
\begin{aligned}
S_Q^{(1)} &= -18\mu^{-\epsilon}\xi_1^2\int d^4\theta_1 d^4\theta_2\int\frac{d^dp}{(2\pi)^d}\partial_{\beta\dot{\beta}}\bar{\sigma}(-p,\theta_1)\partial_{\alpha\dot{\alpha}}\sigma(p,\theta_2)\times\\
&\times\int\frac{d^dk}{(2\pi)^d}\frac{\bar{D}^{\dot{\beta}}D^{\alpha}\bar{D}^2D^2}{16}\delta_{12}\frac{D^{\beta}\bar{D}^{\dot{\alpha}}D^2\bar{D}^2}{16}\delta_{12}G_2(k)G_2(k+p).
\end{aligned}
\tag{10.14}
$$

Carrying out the transformations analogous to those used above we obtain

$$
S_Q^{(1)} = -\mu^{-\epsilon}\int d^4\theta d^dx\partial_{\alpha\dot{\alpha}}\sigma\partial^{\alpha\dot{\alpha}}\bar{\sigma}(\frac{18\xi_1^2(4\pi)^4}{Q^4\epsilon}+fin)\equiv S^{(1)}_{Q_{div}} + S^{(1)}_{Q_{fin}}.
\tag{10.15}
$$

After introducing the one-loop counterterm $-S^{(1)}_{Q\ div}$ one will obtain using (10.12)

$$Z_Q = (1 + \frac{32\pi^2}{Q^2}\frac{18\xi_1^2(4\pi)^4}{Q^4\epsilon}). \qquad (10.16)$$

So we have studied renormalization of ξ_1, ξ_2 and Q^2. As for the Λ, it was already noted that all diagrams containing vertex of type V_4 are finite, it means that the coupling Λ is not renormalized.

Now it remains to investigate renormalization of m^2. It follows from (10.9), that divergent diagrams can contain no more than one vertex of V_3-type corresponding to coupling constant m^2. All other possible vertices should be of V_1- or V_2-types.

We will study the divergent corrections to m^2 in the case when $\xi_1 = \xi_2 = 0$. It means that the vertices V_1 and V_2 are absent at all. It will be proved further that this case corresponds to infrared limit of the theory. It means that only V_3-type vertex can be presented in the diagrams giving contribution to divergent correction to m^2. All these diagrams contain only one vertex V_3-type, one internal line G_{+-}-type and an arbitrary number of external lines corresponding to σ, $\bar{\sigma}$



Fig.12

Let us consider such a diagram with a given number N of external lines, l from those are chiral and other are antichiral. Contribution of this diagram has the form

$$-\frac{m^2}{2}\int d^8z \frac{\sigma^l(z)\bar{\sigma}^{N-l}(z)}{l!(N-l)!}G_{+-}(z,z). \qquad (10.17)$$

Sum of all these contributions is equal to

$$S_3 = -\frac{m^2}{2}\int d^8z \sum_{N=0}^{\infty}\sum_{l=0}^{N}\frac{\sigma^l(z)\bar{\sigma}^{N-l}(z)}{l!(N-l)!}G_{+-}(z,z) = \frac{m^2}{2}\int d^8z e^{\sigma+\bar{\sigma}}G_{+-}(z,z). \qquad (10.18)$$

In momentum representation the S_3 can be written as follows

$$S_3 = -\frac{m^2}{2}\int d^4\theta d^dx e^{\sigma+\bar{\sigma}}\int \frac{d^dk}{(2\pi)^d}\frac{-A^2k^2+m^2}{k^2(-A^2k^2+m^2)^2 - 81\Lambda\bar{\Lambda}}\frac{\bar{D}_1^2D_1^2}{16}\delta_{11}. \qquad (10.19)$$

Taking into account $\frac{\bar{D}_1^2D_1^2}{16}\delta_{11} = \frac{\bar{D}_1^2D_1^2}{16}\delta_{12}|_{\theta_1=\theta_2} = 1$ (note that here the tadpole graph depicted at Fig. 13 has the non-zero contribution!) we get

$$S_3 = -\frac{m^2}{2}\int d^4\theta d^dx e^{\sigma+\bar{\sigma}}\int \frac{d^dk}{(2\pi)^d}\frac{-A^2k^2+m^2}{k^2(-A^2k^2+m^2)^2 - 81\Lambda\bar{\Lambda}}. \qquad (10.20)$$

After integration over momentum we get

$$S_3 = \frac{m^2}{2}\int d^dxd^4\theta e^{\sigma+\bar\sigma}(\frac{2}{16\pi^2A^2\epsilon} + fin).$$

We note that despite these diagrams are tadpole-type, their contribution is not equal to zero unlike Wess-Zumino model.

To cancel the divergence we should introduce a counterterm $-S_{3div}$. It corresponds to mass renormalization

$$\begin{aligned} m_0^2 &= \mu^{-\epsilon}Z_m m^2; \\ Z_{m^2} &= 1 + \frac{2}{16\pi^2A^2\epsilon} = 1 + \frac{2}{Q^2\epsilon} \end{aligned} \tag{10.21}$$

Here m_0^2 is a bare mass and m^2 is a renormalized one.

Next step is consideration of beta functions. As usual, beta function for any renormalized parameter $g(\mu)$ is defined as

$$\beta_g = \mu\frac{dg}{d\mu},$$

where the renormalized parameter is expressed in term of the bare one g_0 which does not depend on μ, $\frac{dg_0}{d\mu} = 0$.

Using (10.13,10.12,10.21) we obtain the following beta functions

$$\beta_{\xi_1} = \frac{72(16\pi)^2}{Q^4}\xi_1\xi_2;\ \beta_{\xi_2} = \frac{72(16\pi)^2}{Q^4}\xi_2^2;\ \beta_{Q^2} = 32\pi^2\frac{18(16\pi)^2\xi_1^2}{Q^4}. \tag{10.22}$$

As a result the equations for running couplings have the form

$$\frac{d\xi_1}{dt} = a\frac{\xi_1\xi_2}{Q^4};\ \frac{d\xi_2}{dt} = a\frac{\xi_2^2}{Q^4};\ \frac{dQ^2}{dt} = b\frac{\xi_1^2}{Q^4}, \tag{10.23}$$

where $a = 2^{11}3^2\pi^2$, $b = 3^2 2^{14}\pi^4$. The solutions of these equations are

$$\begin{aligned} \xi_1(t) &= \frac{\xi_1}{\xi_2}\xi_2(t) \\ Q^2(t) &= Q^2 + 8\pi^2\frac{\xi_1^2}{\xi_2^2}(\xi_2(t) - \xi_2) \\ t &= \frac{1}{2^{11}3^2\pi^2}\Big\{ - [Q^2 - 8\pi^2\frac{\xi_1^2}{\xi_2^2}](\frac{1}{\xi_2(t)} - \frac{1}{\xi_2}) - \\ &- 16\pi^2(\frac{\xi_1}{\xi_2})^2[Q^2 - 8\pi^2\frac{\xi_1^2}{\xi_2^2}]\ln\frac{\xi_2(t)}{\xi_2} + \\ &+ 64\pi^4(\frac{\xi_1}{\xi_2})^4(\xi_2(t) - \xi_2)\Big\}. \end{aligned} \tag{10.24}$$

Let us investigate the behaviour of running couplings $\xi_1(t)$, $\xi_2(t)$ and $Q^2(t)$ in infrared domain when $t \to -\infty$. It is easy to see that in this case $\xi_2(t) \to 0$ and hence $\xi_1(t) \to 0$.

It means that $\xi_1^{(0)} = \xi_2^{(0)} = 0$ is an infrared fixed point. For $Q^2(t)$ we obtain $Q^2(t) \rightarrow Q^2 - 8\pi^2 \frac{\xi_1^2}{\xi_2}$. If we take quantities of initial ξ_1 and ξ_2 so that they correspond to infrared fixed point $\xi_1 = \xi_2 = 0$ one gets $Q^2(t) \rightarrow Q^2$. In particular, only the diagrams given on Fig.4 can contribute to mass renormalization in infrared limit.

To investigate a behaviour of running mass we should use a notion of scaling dimension of superfields. We note that the action of the theory (10.3) at $\xi_1 = \xi_2 = 0$ is invariant under the transformations

$$\delta\sigma = (x^a \partial_a + \frac{1}{2}\theta^\alpha D_\alpha)\sigma + 1; \tag{10.25}$$
$$\delta\bar{\sigma} = (x^a \partial_a + \frac{1}{2}\bar{\theta}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}})\bar{\sigma} + 1$$

Let V be is some function depending on superfields σ, $\bar{\sigma}$ and their derivatives $\partial_a\sigma$, $\partial_a\bar{\sigma}$, $D_\alpha\sigma$, $\bar{D}_{\dot{\alpha}}\bar{\sigma}$,.... We call that V has the scaling dimension Δ if the transformation law of V under transformations (10.25) looks like this

$$\delta V[\sigma, \bar{\sigma}] = (x^a \partial_a + \frac{1}{2}\theta^\alpha D_\alpha + \frac{1}{2}\bar{\theta}_{\dot{\alpha}} \bar{D}^{\dot{\alpha}} + \Delta)V. \tag{10.26}$$

It is easy to see that the superfields σ, $\bar{\sigma}$ have no definite scaling dimension, the derivatives $\partial_a\sigma$, $\partial_a\bar{\sigma}$ have scaling dimension equal to 1, the spinor derivatives $D_\alpha\sigma$,$\bar{D}_{\dot{\alpha}}\bar{\sigma}$ have no definite scaling dimensions. However, the functions e^σ, $e^{\bar{\sigma}}$ have definite scaling dimensions Δ=1.

Let us fulfil the transformations $\sigma \rightarrow \alpha\sigma$, $\bar{\sigma} \rightarrow \alpha\bar{\sigma}$ and $S \rightarrow \frac{1}{\alpha^2}S$ in the action (10.3) at $\xi_1 = \xi_2 = 0$. It leads to the following action depending on arbitrary real parameter α

$$S = \int d^8 z(-\frac{1}{2}\frac{Q^2}{16\pi^2}\bar{\sigma}\Box\sigma - \frac{m^2}{2\alpha^2}e^{\alpha(\sigma+\bar{\sigma})}) + (\frac{\Lambda}{\alpha^2}\int d^6 z e^{3\alpha\sigma} + h.c.).$$

We consider the calculation of the renormalization constant Z_m in the theory (10.27). The only modification in comparison with eq. (10.21) is that we should use the propagator $\alpha^2 G_{+-}$ in supergraph given by Fig.4. The parameter α is resulted here because of expansion of $e^{\alpha(\sigma+\bar{\sigma})}$. It leads immediately to

$$Z_{m^2} = 1 + \frac{2\alpha^2}{Q^2\epsilon}$$

Therefore the equation for running mass will be

$$\frac{dm^2(t)}{dt} = \frac{2\alpha^2 m^2(t)}{Q^2} + \Delta_{m^2} m^2(t). \tag{10.27}$$

where Δ_{m^2} is a scaling dimension of $m^2(t)$.

To find Δ_{m^2} we consider the term $\frac{m^2}{2\alpha^2}e^{\alpha(\sigma+\bar{\sigma})}$ in the action (10.27). The scaling dimension of this term is -2, α is dimensionless and scaling dimension of $e^{\alpha(\sigma+\bar{\sigma})}$ is 2α. Hence $\Delta_{m^2} = 2 - 2\alpha$. Therefore the equation (10.27) looks like this

$$\begin{aligned} \frac{dm^2(t)}{dt} &= (2 - 2\alpha + \frac{2\alpha^2}{Q^2})m^2(t); \qquad (10.28) \\ m^2(0) &= m^2 \end{aligned}$$

where we took into account that $Q^2(t) = Q^2$ in infrared limit. A solution of this equation can be written in the form

$$m^2(t) = m^2 \exp((2 - 2\alpha + \frac{2\alpha^2}{Q^2})t). \tag{10.29}$$

It is evident that at $2 - 2\alpha + \frac{2\alpha^2}{Q^2} > 0$ we get $m^2(t) \to 0$ in infrared limit (note that this condition is satisfied at $\alpha = 1$, i.e. when there is no rescaling). It corresponds to $\kappa^2(t) \to \infty$ where $\kappa^2(t)$ is the running gravitational constant.

As for coupling constant Λ, its beta-function is equal to zero since the vertex of V_4-type is always finite (see above) and the fields σ, $\bar{\sigma}$ are not renormalized in this approach.

Therefore in infrared limit we stay with the following action

$$S = \int d^8 z (-\frac{1}{2}\frac{Q^2}{16\pi^2}\bar{\sigma}\Box\sigma + (\Lambda \int d^6 z e^{3\sigma} + h.c.). \tag{10.30}$$

Our aim consists of calculation of low-energy leading contributions to one-loop effective action. As usual, the first step is background-quantum splitting

$$\sigma \to \sigma + \chi, \bar{\sigma} \to \bar{\sigma} + \bar{\chi}. \tag{10.31}$$

Here $\sigma, \bar{\sigma}$ are background superfields, and $\chi, \bar{\chi}$ are quantum ones. It is known that to find one-loop contribution to effective action it is enough to consider the quadratic action of quantum superfields which has the form

$$S^{(2)}_q[\sigma, \bar{\sigma}; \chi, \bar{\chi}] = \int d^8 z (-\frac{1}{2}\frac{Q^2}{16\pi^2}\bar{\chi}\Box\chi + (\frac{9}{2}\Lambda \int d^6 z e^{3\sigma}\chi^2 + h.c.).$$

The one-loop effective action $\Gamma^{(1)}[\sigma, \bar{\sigma}]$ can be read off from expression

$$\exp(i\Gamma^1[\sigma, \bar{\sigma}]) = \int D\chi D\bar{\chi} \exp(iS^{(2)}_q[\sigma, \bar{\sigma}; \chi, \bar{\chi}]). \tag{10.32}$$

We suggest that the effective action, as usual, has the structure described by (9.2,9.3). The one-loop effective action in this theory can be expressed in the form of effective action of some real scalar superfield just as we done in Wess-Zumino model: we consider the theory of a real superfield v with action

$$S_v = \frac{Q^2}{16\pi^2}\frac{1}{16}\int d^8 z v D^\alpha \bar{D}^2 D_\alpha \Box v. \tag{10.33}$$

One-loop effective action W_v for this theory in the framework of Faddeev-Popov approach is given by

$$e^{iW_v} = \int Dv \exp(iS_v)\delta(\frac{1}{4}D^2 v - \bar{\chi})\delta(\frac{1}{4}\bar{D}^2 v - \chi)\Delta, \tag{10.34}$$

where Δ is Faddeev-Popov determinant which is a constant as in Wess-Zumino model. We note that W_v is also a constant. After multiplying of left-hand side and right-hand side of equations (10.32) and (10.34) respectively and integration over $\chi, \bar{\chi}$ we get one-loop contribution to effective action $\bar{\Gamma}^{(1)}$ in the form

$$\Gamma^{(1)} = -\frac{i}{2}\mathrm{Tr}\log(\frac{Q^2}{16\pi^2}\Box^2 - 9\Lambda e^{3\sigma}\frac{\bar{D}^2}{4} - 9\Lambda e^{3\bar{\sigma}}\frac{D^2}{4}). \tag{10.35}$$

In Schwinger representation the one-loop contribution to effective action looks like

$$\Gamma^{(1)} = -\frac{i}{2}\int_0^\infty \frac{ds}{s}\mathrm{Tr}\exp[is(\frac{Q^2}{16\pi^2}\Box^2 - 9\Lambda e^{3\sigma}\frac{\bar{D}^2}{4} - 9\Lambda e^{3\bar{\sigma}}\frac{D^2}{4})]. \tag{10.36}$$

After change $s\frac{Q^2}{16\pi^2} \to s, \Lambda\frac{16\pi^2}{Q^2} \to \Lambda$ we can express this effective action as

$$\Gamma^{(1)} = -\frac{i}{2}\int_0^\infty \frac{ds}{s}\mathrm{Tr}e^{is\Box^2}\exp[is(-9\Lambda e^{3\sigma}\frac{\bar{D}^2}{4} - 9\Lambda e^{3\bar{\sigma}}\frac{D^2}{4})]. \tag{10.37}$$

Really, commutators of $\Box^2$ with background superfields can lead only to terms depending on space-time derivatives of background superfields which lead only to higher orders in $\partial_a\sigma$. We can calculate exponent of $\tilde{\Delta} = -9\Lambda e^{3\sigma}\frac{\bar{D}^2}{4} - 9\Lambda e^{3\bar{\sigma}}\frac{D^2}{4}$ by the same way as in Wess-Zumino model. The necessary expressions are

$$\begin{aligned} e^{is\Box^2}\delta^4(x_1 - x_2)|_{x_1=x_2} &= \int \frac{d^4k}{(2\pi)^4}e^{isk^4} = \frac{1}{32is\pi^2} && (10.38)\\ \Box e^{is\Box^2}\delta^4(x_1 - x_2)|_{x_1=x_2} &= \int \frac{d^4k}{(2\pi)^4}(-k^2)e^{isk^4} = -\frac{\sqrt{\pi}}{32\pi^2(is)^{3/2}}. && (10.39)\end{aligned}$$

The expression (10.37) can be exactly found in two special cases:

(i) kählerian effective potential, in this case all derivatives of $\sigma, \bar{\sigma}$ are equal to zero. The expression (10.37) is analogous to the expression (8.16) after redefinitions $9\Lambda e^{3\sigma} \to \psi, 9\Lambda e^{3\bar{\sigma}} \to \bar{\psi}$. As a result we can easily restore expression for Schwinger coefficients $A(s), \tilde{A}(s)$ (6.9):

$$A(s) = \tilde{A}(s) = \frac{1}{\Box}[\cosh(9is\Lambda\sqrt{e^{3(\sigma+\bar{\sigma})}\Box}) - 1]. \tag{10.40}$$

The one-loop kählerian effective potential is given by

$$K^{(1)} = -\frac{i}{2}\int_0^\infty [A(s) + \tilde{A}(s)]U(x, x'; s)|_{x=x'}, \tag{10.41}$$

where $U(x, x'; s) = e^{is\Box^2}\delta^4(x - x')$. We can write

$$K^{(1)} = -\frac{i}{2}\int_0^\infty \frac{d\tilde{s}}{\tilde{s}}\sum_{n=0}^\infty \Big[\frac{(9\Lambda\tilde{s}e^{\frac{3}{2}(\sigma+\bar{\sigma})})^{2n+2}}{(2n+2)!}\Box^n\Big]e^{is\Box^2}\delta^4(x - x')|_{x=x'}. \tag{10.42}$$

We can separate sum over n into sum over odd n and sum over even n. We use expressions (10.38) and take into account that

$$\begin{aligned} \Box^{2n}e^{is\Box^2}\delta^4(x_1 - x_2) &= (\frac{\partial}{\partial is})^n e^{is\Box^2}\delta^4(x_1 - x_2) && (10.43)\\ \Box^{2n+1}e^{is\Box^2}\delta^4(x_1 - x_2) &= (\frac{\partial}{\partial is})^n \Box e^{is\Box^2}\delta^4(x_1 - x_2). && (10.44)\end{aligned}$$

As a result we find one-loop kählerian effective potential in the form

$$K^{(1)} = c\Lambda^{2/3}(\frac{16\pi^2}{Q^2})^{2/3}e^{\sigma+\bar{\sigma}}, \tag{10.45}$$

where c is a constant given in [38].

(ii) chiral effective Lagrangian. To calculate it we put $\bar{\sigma} = 0$ which leads to calculation of

$$\tilde{U}(\sigma|s) = \exp[is(-9\Lambda e^{3\sigma})\frac{\bar{D}^2}{4} - 9\Lambda\frac{D^2}{4})] \tag{10.46}$$

up to the second (leading) order is spinor derivatives of σ. We note that as a result one-loop effective action in leading order takes the form

$$\begin{aligned} \Gamma^{(1)} &= \frac{i}{2}\int d^8z \int_0^\infty \frac{ds}{s}[A_1(\sigma|\Box)D^\alpha\sigma D_\alpha\sigma + A_2(\sigma|\Box)D^2\sigma + A_3(\sigma|\Box)] \times \\ &\times \; U_0(x, x'; s)|_{x=x'}. \end{aligned} \tag{10.47}$$

After transforming of this expression to the form of integral over d^6z it is at least of second order in space-time derivatives of σ. Hence we can put all coefficients A_1, A_2, A_3 to depend only on σ but not on its derivatives. It means that at the step of calculating $\tilde{U}(\sigma|s)$ we can put all space-time derivatives of $\sigma, \bar{\sigma}$ to zero. After calculation of $\tilde{U}(\sigma|s)$ (10.46) we get the coefficients of Schwinger expansion $A(s), \tilde{A}(s)$ (note again that only they contribute to one-loop effective action, see (10.41)):

$$\tilde{A}(s) = \frac{1}{\Box}[\cosh(iWs) - 1] \tag{10.48}$$

$$\begin{aligned} A(s) &= -16h(-8h\phi(is)^2 + 64\frac{h^2}{W^2}(\frac{\sinh(iWs)}{W} - is)D^2\phi + \\ &+ \; 8192\frac{h^4}{W^5}[iWs\cosh(iWs) - 3\sinh(iWs) + 2iWs]D^\alpha\phi D_\alpha\phi. \end{aligned} \tag{10.49}$$

Here $\phi = e^{3\sigma}$, $h = -\frac{9}{4}\Lambda\frac{16\pi^2}{Q^2}$, $W = 16he^{3/2\sigma}\sqrt{\Box}$. This expression for A and $\tilde{A}$ can be substituted into (10.47). After expansion of A and $\tilde{A}$ into power series, transformation of the contribution to the form of integral over d^6z and integration over s (see details in [38]), one-loop leading chiral contribution to effective action takes the form

$$\mathcal{L}_c^{(1)} = \Lambda^{1/3}\Big[\{(c_1 + 3c_3)e^{-\sigma} + c_2e^{2\sigma} + 3c_4e^{-4\sigma}\}\partial^m\sigma\partial_m\sigma + (c_3e^{-\sigma} + c_4e^{-4\sigma})\Box\sigma\Big]. \tag{10.50}$$

Here c_1, c_2, c_3, c_4 are finite constants given in [38]. We find that the leaing chiral contribution to effective action is of second order in space-time derivatives of chiral superfield σ, therefore one-loop chiral effective potential is absent.

Then, if we sum classical action (10.3) and leading quantum corrections (10.45,10.50) we get one-loop corrected effective action. If we put in it all derivatives of superfields to zero we get the following low-energy leading effective action:

$$\Gamma = c(\frac{\Lambda}{Q^2})^{2/3}\int d^8z e^{\sigma+\bar{\sigma}} + (\Lambda\int d^6z e^{3\sigma} + h.c.). \tag{10.51}$$

We remind that $\sigma = \log\Phi$, $\bar{\sigma} = \log\bar{\Phi}$ where $\Phi, \bar{\Phi}$ are chiral and antichiral supergravity prepotentials (so called chiral compensators). Expression of this effective action in terms of $\Phi, \bar{\Phi}$ gives

$$\Gamma = c(\frac{\Lambda}{Q^2})^{2/3}\int d^8z\Phi\bar{\Phi} + (\Lambda\int d^6z\Phi^3 + h.c.). \tag{10.52}$$

This action has the structure similar to the classical action of Wess-Zumino model. Hence we see that Wess-Zumino model is generated at infrared limit of four-dimensional dilaton supergravity.

11 Supergauge theories

This section is a brief review of results on supergauge theories. Unfortunately, restricted volume of this section does not allow to discuss all essential results of last years in this sphere hence we only give here main ones.

The starting point of our consideration is an action of $\mathcal{N} = 1$ super-Yang-Mills theory:

$$S_{SYM} = \frac{1}{4g^2}\int d^6z \mathrm{tr} W^\alpha W_\alpha, \tag{11.1}$$

where

$$W_\alpha = -\frac{1}{8}\bar{D}^2(e^{-2gV}D_\alpha e^{2gV}); V(z) = V^I(z)T^I. \tag{11.2}$$

The $V(z) = V^I(z)T^I$ is a real scalar Lie-algebra-valued superfield. We can expand the action (11.1) into power series in coupling g. As a result we get

$$S = \frac{1}{16}\int d^8z \mathrm{tr}(VD^\alpha\bar{D}^2D_\alpha V + \ldots). \tag{11.3}$$

Here dots denote higher orders in g. The action (11.1) is invariant under gauge transformations

$$e^{2gV} \to e^{-2ig\bar{\Lambda}}e^{2gV}e^{2ig\Lambda}. \tag{11.4}$$

where $\bar{D}_{\dot{\alpha}}\Lambda = 0$. The equivalent form of this transformation [10] is

$$\delta(gV) = L_{gV}(2ig\Lambda - 2ig\bar{\Lambda} + \coth_{gV}(2ig\Lambda + 2ig\bar{\Lambda})). \tag{11.5}$$

Here $L_{gV}A = [gV, A]$ is a Lie derivative. It is easy to see that strengths $W_\alpha, \bar{W}_{\dot{\alpha}}$ are invariant under such transformations. The leading order in (11.5) is

$$\delta(gV) = 2ig(\Lambda - \bar{\Lambda}). \tag{11.6}$$

Since the theory is gauge invariant we must introduce gauge-fixing functions for quantization. The most natural form of them is (cf. section 8 where these gauge-fixing functions were used for calculation of one-loop effective action in Wess-Zumino model)

$$\begin{aligned} \chi(V) &= -\frac{1}{4}\bar{D}^2 V + f(z) \\ \bar{\chi}(V) &= -\frac{1}{4}D^2 V + \bar{f}(z). \end{aligned} \tag{11.7}$$

Here $f(z)$ is arbitrary chiral superfield. Variation of these gauge fixing functions under transformations (11.5) is

$$\delta \begin{pmatrix} \chi(V) \\ \bar{\chi}(V) \end{pmatrix} = \begin{pmatrix} 0 & -\frac{1}{4}\bar{D}^2 \\ -\frac{1}{4}D^2 & 0 \end{pmatrix} \begin{pmatrix} g\Lambda \\ g\bar{\Lambda} \end{pmatrix}. \tag{11.8}$$

According to Faddeev-Popov approach we can introduce the ghost action

$$S_{GH} = \mathbf{c}'\delta\chi|_{g\Lambda=c, g\bar{\Lambda}=\bar{c}} \tag{11.9}$$

i.e. parameters of transformation $g\Lambda, g\bar{\Lambda}$ are ghosts. Here $\delta\chi \equiv \delta \begin{pmatrix} \chi(V) \\ \bar{\chi}(V) \end{pmatrix}$. from (11.8), $\mathbf{c}'$ is a line $(c'\bar{c}')$ and since Λ is chiral c, c' are also chiral ones. Here c, c' are chiral ghosts and $\bar{c}, \bar{c}'$ are antichiral ones. As usual, ghosts are fermions.

Therefore

$$S_{GH} = \int d^6 z \mathrm{tr} c' \frac{\delta\chi}{\delta V}\delta V + \int d^6 \bar{z} \mathrm{tr} \bar{c}' \frac{\delta\bar{\chi}}{\delta V}\delta V, \tag{11.10}$$

where

$$\delta V = L_{gV}(c - \bar{c} + \coth_{gV}(c + \bar{c})). \tag{11.11}$$

Therefore the action of ghosts looks like

$$S_{GH} = \int d^8 z \mathrm{tr}(c' + \bar{c}') L_{gV}(c - \bar{c} + \coth_{gV}(c + \bar{c})). \tag{11.12}$$

Then, the generating functional for this theory at zero sources according to Faddeev-Popov approach looks like

$$Z[J]|_{J=0} = \int DvD\{c\} e^{i(S_{SYM} + S_{GH})} \delta_+(\frac{1}{4}\bar{D}^2 V - f)\delta_-(\frac{1}{4}D^2 V - \bar{f}). \tag{11.13}$$

Here $D\{c\} \equiv DcDc'D\bar{c}D\bar{c}'$ We can average over functions f and $\bar{f}$ with weight

$$\exp(\frac{i}{\xi}\int d^8 z (f\bar{f} + b\bar{b})), \tag{11.14}$$

where ξ is a some number (gauge parameter). The $b, \bar{b}$ are Nielsen-Kallosh ghosts (in this case their contribution to effective action is a constant, but in background-covariant formulation it is non-trivial). As a result (11.13) takes the form

$$Z[J]_{J=0} = \int DvD\{c\} e^{i(S_{SYM} + S_{GH} + S_{GF})}, \tag{11.15}$$

where

$$S_{GF} = \frac{1}{16\xi} \int d^8 z \mathrm{tr} (\bar{D}^2 V)(D^2 V) \tag{11.16}$$

is a gauge-fixing action [10].

We introduce the total action

$$S_{total} = S_{SYM} + S_{GF} + S_{GH} \tag{11.17}$$

and the generating functional

$$\begin{aligned} Z[J, \{\eta\}] &= \int DV D\{c\} \exp(i(S_{total} + \int d^8 z \mathrm{tr} JV + \int d^6 z \mathrm{tr}(\eta' c' + \eta c) + \\ &+ \int d^6 \bar{z} (\bar{\eta}' \bar{c}' + \bar{\eta} \bar{c}))). \end{aligned} \tag{11.18}$$

Here $\{\eta\}$ is the set of all sources: $\eta, \eta', \bar{\eta}, \bar{\eta}'$. To develop diagram technique we must split action S_{total} into free (quadratic) part and vertices. It is easy to see ([10, 39]) that

$$e^{-2gV} D^\alpha e^{2gV} = 2g D_\alpha V - 2g^2 [V, D_\alpha V] + \frac{4}{3} [V, [V, D_\alpha V]] + \dots. \tag{11.19}$$

Therefore (11.1) looks like

$$\begin{aligned} S_{SYM} &= \int d^8 z \mathrm{tr} \Big(\frac{1}{16} V D^\alpha \bar{D}^2 D_\alpha V + \frac{1}{8} g (\bar{D}^2 D^\alpha V)[V, D_\alpha V] - \\ &- \frac{1}{16} g^2 [V, D^\alpha V] \bar{D}^2 [V, D_\alpha V] - \frac{1}{12} g^2 (\bar{D}^2 D^\alpha V)[V, [V, D_\alpha V]] + \dots \Big). \end{aligned} \tag{11.20}$$

And the ghost action is

$$S_{GH} = \int d^8 z \mathrm{tr} \Big(\bar{c}' c - \bar{c} c' + g (c' + \bar{c}')[V, c - \bar{c}] + \frac{g^2}{3} (c' + \bar{c}')[V, [V, c + \bar{c}]] \Big) + \dots. \tag{11.21}$$

This expression is enough in one- and two-loop calculations.

The quadratic action is

$$S_0 = \frac{1}{2} \int d^8 z \mathrm{tr} V \Big(- \Box + \frac{1}{16} (1 + \frac{1}{\chi} \{D^2, \bar{D}^2\}) \Big) V + \int d^8 z \mathrm{tr} (\bar{c}' c - \bar{c} c'). \tag{11.22}$$

Vertices can be read off from (11.20,11.21). Propagators look like

$$\begin{aligned} < V(z_1) V(z_2) > &= - \frac{1}{\Box} \Big(- \frac{1}{8\Box} D^\alpha \bar{D}^2 D_\alpha + \xi \frac{\{D^2, \bar{D}^2\}}{16\Box} \Big) \delta^8 (z_1 - z_2) \qquad (11.23) \\ < \bar{c}'(z_1) c(z_2) > &= < c'(z_1) \bar{c}(z_2) > = \frac{1}{\Box} \delta^8 (z_1 - z_2). \end{aligned}$$

We note that ghosts are fermions hence any ghost loop corresponds to minus sign. Then, D-factors are associated with vertices containing ghosts just by the same rule as with vertices containing any chiral superfields. We note that if we choose $\xi = -1$ (Feynman gauge) the propagator of gauge superfield takes the simplest form

$$< v(z_1)v(z_2) >= \frac{1}{\Box}\delta^8(z_1 - z_2). \tag{11.24}$$

Note that its sign is opposite to the the sign of propagator of chiral superfield.

If we want to introduce interaction of a chiral superfield with the gauge one the quadratic part in $\Phi, \bar{\Phi}$ looks like

$$S = \int d^8z \bar{\Phi}_i (e^{gV})^i_j \Phi^j, \tag{11.25}$$

if chiral superfield Φ_i is transformed under some representation of the gauge group (i.e. it is an isospinor) or

$$S = \int d^8z \mathrm{tr}(\bar{\Phi} e^{-gV} \Phi e^{gV}), \tag{11.26}$$

if chiral superfield $\Phi = \Phi^a T^a$ is Lie-algebra-valued. Note that under gauge transformations (11.4) the chiral superfield is transformed as

$$\Phi \to e^{-2ig\Lambda}\Phi \tag{11.27}$$

for isospinor chiral superfield and as

$$\Phi \to e^{-2ig\Lambda}\Phi e^{2ig\Lambda} \tag{11.28}$$

for Lie-algebra-valued chiral superfield. Note that $\Lambda, \bar{\Lambda}$ are Lie-algebra-valued in both cases. The vertices can be easily obtained by expanding into power series expressions corresponding to interaction: in first case

$$\int d^8z [\bar{\Phi}_i (e^{gV})^i_j \Phi^j - \Phi_i \bar{\Phi}^i] = \int d^8z \sum_{n=1}^{\infty} \frac{1}{n!} \bar{\Phi}_i (gV^n)^i_j \Phi^j, \tag{11.29}$$

and in second one –

$$\int d^8z (\mathrm{tr}(\bar{\Phi} e^{-gV} \Phi e^{gV}) - \bar{\Phi}\Phi) = \int d^8z \mathrm{tr}(\Phi[V, \bar{\Phi}] + \bar{\Phi}[V, \Phi] + \frac{1}{2}[\Phi, [V, [V, \bar{\Phi}]] + \ldots). \tag{11.30}$$

The diagram technique derived now is very suitable for calculations in sector of background $\Phi, \bar{\Phi}$ only and for calculation of divergences.

Let us consider an example. The $N = 2$ super-Yang-Mills theory with matter is described by the action (see f.e. [40]):

$$\begin{aligned} S &= \mathrm{tr}\frac{1}{g^2} \int d^6z W^\alpha W_\alpha + \mathrm{tr} \int d^8z \bar{\Phi} e^{-gV} \Phi e^{gV} + \\ &+ \sum_{i=1}^{n} \Big[ig(\int d^6z Q_i \Phi \tilde{Q}_i + h.c.) + \int d^8z \bar{\tilde{Q}}_i e^{-gV} \tilde{Q}_i + \int d^8z \bar{Q}_i e^{gV} Q_i \Big]. \end{aligned} \tag{11.31}$$

Here Φ is Lie-algebra-valued chiral superfield, and $Q^i, \tilde{Q}_i$ are chiral superfields transformed under mutually conjugated representations of Lie algebra. They are often called matter hypermultiplets. Let us consider the structure of one-loop divergences in this theory. For simplicity we choose Feynman gauge $\xi = -1$ in which the propagator has the most simple structure (11.24), therefore all tadpole diagrams given in [39] evidently vanish.

First we consider contributions to wave function renormalization of Φ field





Here the thin line is propagator of Φ, the thick one – of hypermultiplets $Q, \tilde{Q}$, the wavy one – of real superfield v, the dashed one – from ghosts.

One-loop divergent contributions from these supergraphs are respectively (see f.e. [40])

$$2\sum_M \int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}\Phi^a\bar{\Phi}^b \mathrm{tr}_M(T^aT^b) \tag{11.32}$$

and

$$-\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}\Phi^a\bar{\Phi}^b \mathrm{tr}_{ad}(T^aT^cT^d)\mathrm{tr}_{ad}(T^bT^cT^d) \tag{11.33}$$

Here tr_M denotes trace in representation under which hypermultiplets are transformed. Coefficient 2 is caused by presence of two chiral hypermultiplets Q and $\tilde{Q}$. We see that if $\sum_M \mathrm{tr}_M(T^aT^b) = \mathrm{tr}_{ad}(T^aT^cT^d)\mathrm{tr}_{ad}(T^bT^cT^d)$ there is no divergent contributions to wave function renormalization.

Corrections to hypermultiplet wave function look like





One-loop divergent contributions from these supergraphs are respectively [40]

$$-\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}\bar{Q}_iQ_l(T^a)^{ij}(T^a)^{jl} \tag{11.34}$$

and

$$\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}\bar{Q}_iQ_l(T^a)^{ij}(T^a)^{jl} \tag{11.35}$$

These corrections evidently cancel each other, hence there is no renormalization of hypermultiplet wave function. In both these cases cancellation is caused by the difference in signs of propagators of gauge superfield and chiral superfields.

One-loop contributions to wave function renormalization for gauge superfields are









Contribution of these four supergraphs are respectively given by (see f.e. [41, 40])

$$\begin{aligned}
&\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}V^aV^b\mathrm{tr}_{ad}(T^aT^cT^d)\mathrm{tr}_{ad}(T^bT^cT^d)\\
&\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}V^aV^b\mathrm{tr}_{ad}(T^aT^cT^d)\mathrm{tr}_{ad}(T^bT^cT^d)\\
2\sum_M &\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}V^aV^b\mathrm{tr}_M(T^aT^b)\\
-4 &\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2(k+p)^2}V^aV^b\mathrm{tr}_M(T^aT^b).
\end{aligned} \tag{11.36}$$

Hence if $2\sum_M \mathrm{tr}_M(T^aT^b)+2\mathrm{tr}_{ad}(T^aT^cT^d)\mathrm{tr}_{ad}(T^bT^cT^d)=4\mathrm{tr}_{ad}(T^aT^b)$ (or, as is the same with taking into account condition for $<\phi\bar{\phi}>$-propagator, $\mathrm{tr}_{ad}(T^aT^cT^d)\mathrm{tr}_{ad}(T^bT^cT^d)=\mathrm{tr}_{ad}(T^aT^b)$) these contributions are cancelled and there is no divergent correction to $<VV>$-propagator, and the theory is finite. This mechanism of vanishing divergences is discussed, f.e. in [41], [42], where it is shown to be caused by the $N=2$ superconformal symmetry. The most important example of such theories is $N=4$ super-Yang-Mills theory where we have one pair of hypermultiplets $Q,\tilde{Q}$ and they are transformed under the adjoint representation of Lie algebra. This theory is known to be finite (see f.e. [39, 41]).

However, the approach developed until this place is very useful for consideration of divergences and corrections in the sector of chiral superfields $\Phi, Q, \tilde{Q}$ only. To study contributions depending on gauge superfields we must develop a method allowing to preserve manifest gauge invariance at any step as earlier we obtained contributions in terms of superfield V which are in general case not gauge invariant. Therefore we must introduce an approach in which external lines are background strengths $W_\alpha, \bar{W}_{\dot{\alpha}}$ and their **gauge covariant** derivatives. This method was developed in [28] (see also [44] and references therein), here we give its description.

The problem of calculation of the effective action in the theory described by the action (11.1) is much more complicated than in other field theories. The main difficulties are the following ones. First, the nonpolynomiality of the action (11.1) implies in the infinite number of vertices which seems to result in infinite number of types of the divergent quantum corrections (such a situation is treated in common cases as the non-renormalizability of the theory), second, it is easy to see that the common background-quantum splitting $V \to V_0 + v$ where V_0 is a background field and v is a quantum field cannot provide manifest gauge covariance of the quantum corrections. Really, because of the nonpolynomiality of W^α (11.2), to get a covariant quantum correction (which by definition must be expressed in terms of the strengths $W_\alpha, \bar{W}_{\dot\alpha}$ which are the only objects transforming covariantly under the gauge transformations unlike of the superfield V itself) we need to summarize an infinite number of supergraphs with different numbers of external V_0 legs. The background field method provides an effective solution for both these problems.

The starting point of the method under discussion is a **nonlinear** background-quantum splitting for the superfield V defining the action (11.1) [28]:

$$e^{2gV} \to e^{2g\Omega} e^{2gv} e^{2g\bar\Omega}. \tag{11.37}$$

Here the v is a quantum field, the $\Omega, \bar\Omega$ are the background superfields (they are not necessary chiral/antichiral ones). After such a background-quantum splitting, the classical action (11.1) takes the form:

$$S = -\frac{1}{64g^2}\int d^8z (e^{-2gv}\mathcal{D}^\alpha e^{2gv})\bar{\mathcal{D}}^2(e^{-2gv}\mathcal{D}_\alpha e^{2gv}), \tag{11.38}$$

In this expression (which describes a theory of real scalar superfield v coupled to background superfields $\Omega, \bar\Omega$) the $\mathcal{D}^\alpha, \bar{\mathcal{D}}^{\dot\alpha}$ are the background covariant derivatives defined by the expressions [41, 28]:

$$\begin{aligned} \mathcal{D}^\alpha &= e^{-2g\Omega} D^\alpha e^{2g\Omega}, \\ \bar{\mathcal{D}}^{\dot\alpha} &= e^{2g\bar\Omega} \bar{D}^{\dot\alpha} e^{-2g\bar\Omega}. \end{aligned} \tag{11.39}$$

Our further aim consists in study of the action (11.38). To do it let us first describe the properties of the background covariant derivatives given by (11.39). The $D^\alpha, \bar{D}^{\dot\alpha}$ in (11.39) act on all on the right. Then, as well as the covariant derivatives in usual differential geometry, the background covariant derivatives $\mathcal{D}^\alpha, \bar{\mathcal{D}}^{\dot\alpha}$ can be represented in the following "standard" form

$$\mathcal{D}^\alpha = D^\alpha - i\Gamma^\alpha, \quad \bar{\mathcal{D}}^{\dot\alpha} = \bar{D}^{\dot\alpha} - i\bar\Gamma^{\dot\alpha}, \tag{11.40}$$

where

$$\Gamma^\alpha = ie^{-2g\Omega}(D^\alpha e^{2g\Omega}), \quad \bar\Gamma^{\dot\alpha} = ie^{2g\bar\Omega}(\bar{D}^{\dot\alpha} e^{-2g\bar\Omega}) \tag{11.41}$$

are the superfield connections.

Let us study the (anti)commutation relations for the $\mathcal{D}^\alpha, \bar{\mathcal{D}}^{\dot\alpha}$. We start with imposing the following constraint

$$\mathcal{D}_{\alpha\dot\alpha} = -\frac{i}{2}\{\mathcal{D}_\alpha, \bar{\mathcal{D}}_{\dot\alpha}\} \tag{11.42}$$

which represent itself as a background covariant analog of the common anticommutation relation $\partial_{\alpha\dot{\alpha}} = -\frac{i}{2}\{D_\alpha, \bar{D}_{\dot{\alpha}}\}$. Then, it is easy to verify straightforwardly the following definition of the background strength W_α (cf. [43]):

$$W_\alpha = -\frac{i}{\sqrt{2}}[\bar{\mathcal{D}}^{\dot{\alpha}}, \{\bar{\mathcal{D}}_{\dot{\alpha}}, \mathcal{D}_\alpha\}] = \sqrt{2}[\bar{\mathcal{D}}^{\dot{\alpha}}, \mathcal{D}_{\alpha\dot{\alpha}}]. \tag{11.43}$$

Really, after we substitute expressions (11.39) for the background-covariant derivatives to (11.43) and take into account that $e^{2g\Omega}e^{2g\bar{\Omega}} = e^{2gV}$ in the case of absence of the quantum field v (cf. (11.37)) we get just the definition (11.2).

The relation (11.43) is crucial. Its treating consists of the fact that the background-covariant space-time derivative $\mathcal{D}_{\alpha\dot{\alpha}}$ has non-zero commutators with spinor background-covariant derivatives unlike of the common covariant derivatives, and, moreover, that the background strengths could arise during the D-algebra transformations. In part, the identity (11.43) implies in the following expressions [44]:

$$\begin{aligned}
[\mathcal{D}_\alpha, \bar{\mathcal{D}}^2] &= 4i(-\sqrt{2}W^\alpha + \bar{\mathcal{D}}^{\dot{\alpha}}\mathcal{D}_{\alpha\dot{\alpha}}) = \\
&= 4i(\sqrt{2}W^\alpha + \mathcal{D}_{\alpha\dot{\alpha}}\bar{\mathcal{D}}^{\dot{\alpha}}).
\end{aligned} \tag{11.44}$$

Another consequence of (11.43), and consequently of (11.44), is

$$-\frac{1}{8}\mathcal{D}^\alpha\bar{\mathcal{D}}^2\mathcal{D}_\alpha + \frac{1}{16}\{\mathcal{D}^2, \bar{\mathcal{D}}^2\} = \Box - \frac{i}{2}W^\alpha\mathcal{D}_\alpha - \frac{i}{2}\bar{W}^{\dot{\alpha}}\mathcal{D}_{\dot{\alpha}}. \tag{11.45}$$

This expression is a gauge covariant analog of the known relation
$-\frac{1}{8}D^\alpha\bar{D}^2D_\alpha + \frac{1}{16}\{D^2, \bar{D}^2\} = \Box$.

Basing on (11.43), we also can prove the following important relations:

$$\begin{aligned}
\mathcal{D}^2\bar{\mathcal{D}}^2\mathcal{D}^2 &= 16(\Box - \frac{i}{2}\bar{W}^{\dot{\alpha}}\bar{\mathcal{D}}_{\dot{\alpha}} - \frac{i}{4}(\bar{\mathcal{D}}^{\dot{\alpha}}\bar{W}_{\dot{\alpha}}))\mathcal{D}^2 \equiv 16\Box_-\mathcal{D}^2; \\
\bar{\mathcal{D}}^2\mathcal{D}^2\bar{\mathcal{D}}^2 &= 16(\Box - \frac{i}{2}W^\alpha\mathcal{D}_\alpha - \frac{i}{4}(\mathcal{D}^\alpha W_\alpha))\bar{\mathcal{D}}^2 \equiv 16\Box_-\bar{\mathcal{D}}^2.
\end{aligned} \tag{11.46}$$

We also need in definition of the (background) covariantly chiral superfields to describe coupling of the gauge superfields to the matter. By definition, the superfield Φ is referred as the (background) covariantly chiral one if it satisfies the condition $\bar{\mathcal{D}}_{\dot{\alpha}}\Phi = 0$. It is easy to see that the Φ is related to the common chiral field Φ_0 as

$$\Phi = e^{g\bar{\Omega}}\Phi_0. \tag{11.47}$$

Really, condition of chirality $\bar{D}_{\dot{\alpha}}\Phi_0 = 0$ implies in

$$e^{g\bar{\Omega}}\bar{D}_{\dot{\alpha}}e^{-g\bar{\Omega}}e^{g\bar{\Omega}}\Phi_0 = 0. \tag{11.48}$$

Using the definitions (11.39,11.47) we arrive just to the condition $\bar{\mathcal{D}}_{\dot{\alpha}}\Phi = 0$.

Now we are in position to develop the petrurbative approach for the theory with action (11.38). First, we note that this theory possesses the symmetry with respect to the following gauge transformations:

$$e^{gv} \rightarrow e^{ig\bar{\Lambda}}e^{gv}e^{-ig\Lambda} \tag{11.49}$$

where Λ is a **covariantly** chiral parameter, i.e. it satisfies the condition $\bar{\mathcal{D}}_{\dot{\alpha}}\Lambda = 0$ (similarly, $\mathcal{D}_{\alpha}\bar{\Lambda} = 0$). Therefore we need to introduce a gauge fixing. The most natural background covariant gauge fixing term looks like

$$S_{gf} = -\frac{1}{16}\int d^8 z v\{\mathcal{D}^2, \bar{\mathcal{D}}^2\}v \tag{11.50}$$

which is a covariant generalization of the common gauge fixing term in the Feynman gauge. Summarizing the (11.38) and (11.50) and taking into account (11.45) we get the following action of the quantum v field:

$$S_t = S + S_{gf} = -\frac{1}{2}\int d^8 z v \Box v + S_{int}. \tag{11.51}$$

In principle, one can fix other gauges (by putting the factor ξ^{-1} in S_{gf}), however, even problem of finding the propagator for v field appears to be very complicated (up to this time nobody found this propagator in a closed form). Hence the Feynman gauge is the most convenient one. The propagator of the v field in the Feynman gauge is

$$< v(z_1)v(z_2) >= \Box^{-1}\delta^8(z_1 - z_2) \tag{11.52}$$

(for the sake of the uniqueness of the consideration we relate all terms involving $W_{\alpha}, \bar{W}_{\dot{\alpha}}$ to the interaction part).

The S_{int} in the expression above is an interaction part. It involves both covariant generalizations of the "common" vertices and the new vertices involving the background strengths $W_{\alpha}, \bar{W}_{\dot{\alpha}}$ manifestly:

$$\begin{aligned} S_{int} &= \int d^8 z \Big(v(-\frac{i}{2}W^{\alpha}\mathcal{D}_{\alpha} - \frac{i}{2}\bar{W}^{\dot{\alpha}}\mathcal{D}_{\dot{\alpha}})v + \\ &+ \frac{1}{8}gv\{\mathcal{D}^{\alpha}v, \bar{\mathcal{D}}^2\mathcal{D}_{\alpha}v\} + \frac{1}{3}[[\mathcal{D}^{\alpha}v, v], v]iW_{\alpha} - \\ &- \frac{1}{16}g^2[v, \mathcal{D}^{\alpha}v]\bar{\mathcal{D}}^2[v, \mathcal{D}_{\alpha}v] - \frac{1}{12}g^2\bar{\mathcal{D}}^2\mathcal{D}^{\alpha}v[v, [v, \mathcal{D}_{\alpha}v]] + \frac{1}{6}[[[\mathcal{D}^{\alpha}v, v], v], v]iW_{\alpha}\Big) + \ldots . \end{aligned} \tag{11.53}$$

Because of the gauge symmetry, one needs to introduce ghosts. Since the form of the gauge transformations and gauge fixing action is very similar to the "common" superfield case with only difference consisting in covariant chirality instead of common one, the action of ghosts in this case also will be analogous to the common case with only difference consisting in the fact of the **covariant** chirality of the ghosts c, c' or **covariant** antichirality of the ghosts $\bar{c}, \bar{c}'$:

$$S_{gh} = \int d^8 z(\bar{c}' + c')L_{gv}(c - \bar{c} + \mathrm{cth} L_{gv}(c + \bar{c})) \tag{11.54}$$

which after expansion in the power series gives:

$$S_{gh} = \int d^8 z(\bar{c}'c - \bar{c}c' + (c' + \bar{c}')[v, c + \bar{c}]) + \frac{1}{12}(c' + \bar{c}')[v, [v, c + \bar{c}]] + \ldots). \tag{11.55}$$

The action of the matter after the background-quantum splitting of the gauge fields takes the form

$$S_m = \int d^8 z \bar{\Phi} e^{gv} \Phi, \tag{11.56}$$

where $\Phi, \bar{\Phi}$ are the **covariantly** chiral and antichiral superfields. The propagators of chiral superfields and ghosts are

$$< \bar{\phi}\phi >= -\frac{1}{16}\mathcal{D}^2 \Box_+^{-1} \bar{\mathcal{D}}^2, \quad < \bar{c}' c >=< \bar{c} c' >== -\frac{1}{16}\mathcal{D}^2 \Box_+^{-1} \bar{\mathcal{D}}^2. \tag{11.57}$$

We note that such propagators can be expanded into power series in $W_\alpha, \bar{W}_{\dot{\alpha}}$, see (11.46). The expressions (11.53,11.55,11.56,11.52,11.57) can be used for constructing the supergraphs. The examples of application of the method for the supergraph calculations can be found in [44].

Now let us give a comparative characteristics for the two methods of superfield calculations – the background field method and the "common" method.

The crucial difference is the following one. In the framework of the "common" method the quadratic and linear divergences could arise for the supergraph with arbitrary any number of the external legs. Really, it is easy to show that the superficial degree of divergence for the "common" supergraph is

$$\omega = 2 - \frac{1}{2} N_D - E_\phi, \tag{11.58}$$

where N_D is a number of spinor supercovariant derivatives associated to the external legs, E_ϕ is a number of external chiral (antichiral) legs. We note that the quadratic and/or linear divergences are possible for any number of external v legs. At the same time, in the framework of the background field method by the construction of the background-quantum splitting the only external lines are the background strengths and/or their covariant derivatives. The superficial degree of divergence in this case can be shown to have the form

$$\omega = 2 - \frac{3}{2} N_W - \frac{1}{2} N_D + \epsilon - E_\phi, \tag{11.59}$$

where N_W is a number of the background strength legs, $\epsilon = 1$ for the chiral (antichiral) contribution (which only possible structure is $\int d^6 z W^2$, $\int d^6 \bar{z} \bar{W}^2$), otherwise $\epsilon = 0$; the N_D is the number of derivatives acting on external $W_\alpha, \bar{W}_{\dot{\alpha}}$ legs (the derivatives presenting in each $W_\alpha, \bar{W}_{\dot{\alpha}}$ by definition must not be taken into account!). We see that in framework of this approach only logarithmic overall divergences are possible, they arise for the terms proportional to W^2, with the quadratic and linear subdivergences (which are important if we make a noncommutative generalization) can arise only in subgraphs which are not associated to the external $W_\alpha, \bar{W}_{\dot{\alpha}}$ legs.

From formal viewpoint such a difference has the following origin. Really, the superfield strength by construction contains three spinor derivatives, hence arising of any superfield strength leg in the framework of "common" formalism decreases by three the number of the D-factors which could be converted to momenta (we note that use of the background field method allows to sum automatically the infinite number of "common" graphs and forbids existence of the supergraphs with the superficial quadratic or linear divergence), as a result, the convergence of the supergraph is improved. It is essential in the noncommutative field theory since it means that the only problems could be generated by subgraphs only as each contribution to the effective action is in worst case only logarithmically divergent. In part, it means that there is no contradiction between the result of Bichl et al. [45] according to which the different contributions to the one-loop two-point function of v field in the $U(1)$ NC SYM theory possess quadratic divergences (only their sum is free of dangerous UV/IR mixing) and the result of Zanon et al. [46] according to which all one-loop contributions to the effective action in the same theory are free of the dangerous UV/IR mixing (notice that the calculations in the last paper were carried out in the framework of the background field method); we should mention also use of the background field method in the papers [46] devoted to study of the one-loop effective action in the noncommutative super-Yang-Mills theories. We note that the background field method allows to preserve the gauge covariance at all steps of calculations.

However, the background field method has one disadvantage – presence of nontrivial commutators of the covariant derivatives makes all calculations extremelly difficult from the technical viewpoint.

12 Conclusions

We considered superfield method in supersymmetric field theory. This method allows to preserve manifest supersymmetry at any step of calculations, and the calculations within it turns to be much more compact than within the framework of the component approach.

We studied several examples of superfield theories and presented in details quantum calculations for them. These examples were Wess-Zumino model, general chiral superfield model, dilaton supergravity and $N = 1$ super-Yang-Mills theory with chiral matter. In these theories we developed supergraph technique, studied general form of superfield effective action and calculated low-energy leading contributions to effective action. It is natural to expect that development of superfield quantum calculations in other superfield models formulated in terms of $N = 1$ superfields including different supergravity models is in principle no more difficult. We also discussed the background field method in the supergrauge theories.

Let us briefly discuss other applications and generalizations of the superfield approach in the quantum field theory. In the last years the following most important ways of applications of superfield supersymmetry were developed.

1. **Studying of theories with extended supersymmetry**. It is known that theories with extended supersymmetry possess better renormalization properties, f.e. as $N = 1$ super-Yang-Mills theory is renormalizable, the $N = 4$ super-Yang-Mills theory is finite. The most important examples of theories with extended supersymmetry are $N = 2$ and $N = 4$ super-Yang-Mills theories. During last years numerous results in studying of these theories were obtained (see f.e. [47, 48, 49] and references therein).

It is natural to expect that the most adequate method for consideration of such theories must possess manifest $N = 2$ supersymmetry. Such a method is a harmonic superspace approach developed in [50],[51]. This method is based on consideration of superfields being the functions of bosonic space-time coordinates x^a, two sets of Grassmann coordinates $\theta^{i\alpha}, \bar{\theta}^{i\dot{\alpha}}$ with $i = 1, 2$ and spherical harmonics $u^{\pm i}$. Introducing of analytic superfield [50] allows to develop formulation in terms of unconstrained $N = 2$ superfield and to avoid arising of component fields with higher spins. The formulations of $N = 2$ and $N = 4$ super-Yang-Mills theories in harmonic superspace is given in [50, 51], background field method for these theories is developed in [52, 53, 54], and examples of quantum calculations are given in [48, 49, 55, 56, 57]. The most important results presented in these papers are calculation of holomorphic action of $N = 2$ matter hypermultiplets in external $N = 2$ gauge superfield, calculation of one-loop nonholomorphic effective potential in $N = 4$ super-Yang-Mills theory and proof of its absence in higher loops, calculation of one-loop effective action for $N = 4$ super-Yang-Mills theory for constant strength tensor F_{ab}, calculation of superconformal anomaly of $N = 2$ matter interacting with $N = 2$ supergravity. During last years other important results achieved in these investigations were calculation of contributions depending on derivatives of $N = 2$ super-Yang-Mills strength $\mathcal{W}$ [58, 59], calculation of contributions depending on background matter hypermultiplet fields [60], and development of quantum approach for $N = 3$ super-Yang-Mills theory [61] (the manifestly $N = 3$ supersymmetric approach based on the harmonic superspace technique for the $N = 3$ supersymmetric theory was given in paper [62]).

2. **Noncommutative supersymmetric theories**. Noncommutative theories have been intensively studied during last years. Concept of space-time noncommutativity was introduced to quantum field theory due to some consequences of D-branes theory [63] and to consideration of quantum theories on very small distances where quantum fluctuations of geometry are essential. Consideration of supersymmetric noncommutative theories is quite natural. During last years some interesting results in studying of noncommutative supersymmetric theories were obtained but they were mostly based on component approach. The first superfield results were calculation of leading ($\sim F^4$) correction to one-loop effective action for $N = 4$ super-Yang-Mills theory [46] and formulation of supergraph technique for noncommutative Wess-Zumino model [64]. Further, the quantum superfield studies for the Wess-Zumino model [65] and four-dimensional superfield QED [66] and super-Yang-MIlls theories [67] were carried out. These theory were shown to be consistent in the sense of absence of the nonintegrable UV/IR infrared divergences. Thus, we can speak about construction of the consistent noncommutative generalizations of the supersymmetric theories of electromagnetic, strong and weak interactions. Therefore, the next most important problem could consist in development of the noncommutative supersymmetric generalization for the last fundamental interaction – the gravitational one. However, this problem is of course extremely difficult (see discussion of the problem f.e. in [68, 69]).

3. **Noncommutative superspace.** One more approach in the superfield quantum theory is based on use of the noncommutative superspace [70]. Within it, the fermionic superspace coordinates form the Clifford algebra instead of the Grassmann algebra, which,

in part, leads to the modified construction of the Moyal product. In the framework of this approach, the generalizations of the Wess-Zumino (see f.e. [71] and reference therein), gauge (see f.e. [72] and reference therein) and general chiral superfield model [73] were studied.

Then, there are a lot of applications of superfields approach to problems of supersymmetric quantum field theory (f.e. to studying of AdS/CFT correspondence which was carried out mostly on base of component approach), and of course consideration of many problems originated from superstrings and branes theory.

As a final conclusion, we can suppose that superfield approach in quantum field theory is a very perspective one, and there are a lot of ways for its development and more applications.

Acknowledgements. Author is grateful to Prof. V. O. Rivelles, Prof. M. O. C. Gomes, Prof. H. O. Girotti and Prof. A. J. da Silva for discussions and joint researches, to Prof. I. L. Buchbinder, Prof. M. Cvetic, Prof. S. M. Kuzenko for collaboration, and to Instituto de Fisica, Universidade de São Paulo and Instituto de Fisica, Universidade Federal do Rio Grande do Sul for hospitality. The work was supported by FAPESP, project No. 00/12671-7.

References

[1] M. Green, J. Schwarz, E. Witten. Theory of superstrings, Cambridge, UK: Univ. Pr., 1987, vol. 1-2.

[2] D. V. Volkov, V. P. Akulov. JETP Lett., 16, 621 (1972); Phys. Lett. B46, 109 (1973).

[3] Y. A. Golfand, E. S. Lichtman. JETP Lett., 13, 323 (1971).

[4] J. Wess, B. Zumino. Nucl. Phys. B70, 139 (1974); Nucl. Phys. B78, 1 (1974).

[5] The Supersymmetric World, ed. by G. Kane and M. Shifman, World Scientific, 2000.

[6] J. Wess, B. Zumino. Phys. Lett. B49, 52 (1974).

[7] A. Salam, J. Strathdee. Nucl. Phys. B76, 477 (1974).

[8] S. Ferrara, J. Wess, B. Zumino. Phys. Lett. B51, 239 (1974).

[9] J. Wess, J. Bagger. Supersymmetry and supergravity. Princeton University Press, Princeton, 1983.

[10] I. L. Buchbinder, S. M. Kuzenko. Ideas and Methods of Supersymmetry and Supergravity. IOP Publishing, Bristol and Philadelphia, 1998.

[11] V. Ogievetsky, L. Mezinchesku. Uspehi Fiz. Nauk (Phys. Sci. Achievments) 117, 637 (1975) (in Russian).

[12] I. L. Buchbinder, A. Yu. Petrov. Yad. Fiz. (Phys. Atom. Nucl.) 63, 9, 2557 (2000).

[13] I. L. Buchbinder, A.Yu. Petrov. Class. Quant. Grav. 13, 2081 (1996).

[14] N. Seiberg, E. Witten. Nucl. Phys. B430, 19 (1994).

[15] P. West. Introduction to supersymmetry and supergravity. World Scientific, 1989.

[16] P. West. Phys. Lett. B258, 375 (1991).

[17] I. Jack, D. R. T. Jones. Problems of supersymmetric regularization, in: Kane G. L. (ed.), Perspectives of supersymmetry, 149-167 (hep-ph/9707278).

[18] J. Collins. Renormalization, UK: Univ. Pr., 1984.

[19] I. Jack, D. R. T. Jones, P. West. Phys. Lett. B258, 382 (1991).

[20] P. West. Phys. Lett. B261, 396 (1991).

[21] I. L. Buchbinder, S. M. Kuzenko, A.Yu. Petrov. Phys. Lett. B321, 372 (1994).

[22] I. L. Buchbinder, S. M. Kuzenko, A. Yu. Petrov. Yad. Fiz. (Phys. Atom. Nucl.), 59, 1, 157 (1996).

[23] I. L. Buchbinder, S. D. Odintsov, I. L. Shapiro. Effective action in quantum gravity. IOP Publishing, Bristol and Philadelphia, 1992.

[24] B.S. De Witt. Dynamical theory of groups and fields. Princeton Univ. Press, Princeton, 1987.

[25] I. L. Buchbinder, S. M. Kuzenko, J. V. Yarevskaya. Nucl. Phys. B411, 665 (1994); Yad. Fiz. (Phys. Atom. Nucl), 56, 5, 202 (1993).

[26] I. N. McArthur, T. D. Gargett. Nucl. Phys. B497, 525 (1997).

[27] S. Coleman, S. Weinberg. Phys. Rev. D7, 1888 (1973).

[28] M. T. Grisaru, M. Rocek, W. Siegel. Nucl. Phys. B159, 429 (1979).

[29] A. Pickering, P. West. Phys. Lett. B353, 54 (1997).

[30] A. Yu. Petrov. Superfield effective action in supersymmetric field theories (PhD thesis), Tomsk, 1997.

[31] S.M. Kuzenko, J. V. Yarevskaya. Yad. Fiz. (Phys. Atom. Nucl.) 56, 5, 195 (1993).

[32] G. Cleaver, M. Cvetič, J.R. Espinosa, L. Everett, and P. Langacker, Nucl. Phys. B525, 3, 1998; Phys. Rev, D59, 115003, 1999; M. Cvetič, L. Everett, and J. Wang. Nucl. Phys. B538, 52 (1999).

[33] A.Yu. Petrov. Effective action in general chiral superfield model, *Preprint* hep-th/0002013.

[34] I. L. Buchbinder, A.Yu. Petrov. Phys.Lett. B461, 209 (1999).

[35] K. Symanzik. Commun. Math. Phys, 34, 7 (1973); T. Appelquist and J. Carrazone, Phys. Rev. D11, 2856 (1975).

[36] I. L. Buchbinder, M. Cvetič, A.Yu. Petrov. Mod.Phys. Lett. A15, 783 (2000); Nucl. Phys. B571, 358 (2000).

[37] I. L. Buchbinder, S. M. Kuzenko. Phys. Lett. B202, 233 (1988).

[38] I. L. Buchbinder, A. Yu. Petrov. Class. Quant. Grav. 14, 21 (1997).

[39] S. Kovacs. A perturbative reanalysis of $N = 4$ supersymmetric Yang-Mills theory, *Preprint* hep-th/9902047.

[40] A. De Giovanni, M. T. Grisaru, D. Zanon. Phys. Lett. B409, 251 (1997).

[41] S. J. Gates, M. T. Grisaru, M. Rocek, W. Siegel. Superspace or One Thousand and One Lectures in Supersymmetry, Benjamin/Cummings, 1983.

[42] P. S. Howe, K. Stelle, P. West. Phys. Lett. B124, 55 (1983).

[43] M. T. Grisaru, W. Siegel, Nucl. Phys. B201, 292 (1982).

[44] M. T. Grisaru, D. Zanon, Phys. Lett. B142, 359 (1984); Nucl. Phys. B252, 578 (1985).

[45] A. A. Bichl, M. Ertl, A. Gerhold, J. M. Grimstrup, H. Grosse, L. Popp, V. Putz, M. Schweda, R. Wulkenhaar. Int. J. Mod. Phys. A19, 4231 (2004).

[46] D. Zanon, Phys. Lett. B502, 265 (2001), Phys. Lett. B504, 101 (2001); A. Santambrogio, D. Zanon, JHEP 0101, 024 (2001); M. Pernici, A. Santambrogio, D. Zanon, Phys. Lett. B504, 131 (2001).

[47] I. L. Buchbinder, S. M. Kuzenko, A.A. Tseytlin. Phys. Rev. D62, 045001 (2000).

[48] I. L. Buchbinder, S. M. Kuzenko. Phys. Lett. B446, 216 (1999); Mod. Phys. Lett. A13, 1623 (1998).

[49] I. L. Buchbinder, A. Yu. Petrov. Phys. Lett. B469, 482 (2000).

[50] A. Galperin, E.Ivanov, S. Kalitzin, V. Ogievetsky, E. Sokachev. Class. Quant. Grav. 1, 469 (1984).

[51] A. Galperin, E. Ivanov, V. Ogievetsky, E. Sokachev. Class. Quant. Grav. 2, 601 (1985); 2, 617 (1985).

[52] E.I. Buchbinder, I. L. Buchbinder, E. A. Ivanov, S. M. Kuzenko, B. A. Ovrut. Phys. Lett. B412, 309 (1997).

[53] E. I. Buchbinder, I. L. Buchbinder, S. M. Kuzenko, B. A. Ovrut. Phys. Lett. B417, 61 (1997).

[54] I. L. Buchbinder, S. M. Kuzenko, B. A. Ovrut. Phys. Lett. B433, 335 (1997).

[55] I. L. Buchbinder, I. B. Samsonov. Mod. Phys. Lett. A14, 2537 (1999).

[56] S. M. Kuzenko, I. N. McArthur. Phys. Lett. B506, 140 (2001).

[57] S. M. Kuzenko, S. Theisen. Class. Quant. Grav. 17, 665 (2000).

[58] I. L. Buchbinder, A. Yu. Petrov, A. A. Tseytlin, Nucl. Phys. B621, 179 (2002).

[59] S. M. Kuzenko, I. N. McArthur, JHEP 0310, 029 (2003); Nucl. Phys. B683, 3 (2004); Nucl. Phys. B697, 89 (2004).

[60] I. L. Buchbinder, E. A. Ivanov, A. Yu. Petrov, Nucl. Phys. B653, 64 (2003).

[61] I. L. Buchbinder, E. A. Ivanov, I. B. Samsonov, B. M. Zupnik, Nucl. Phys. B689, 91 (2004).

[62] F. Delduc, I. McCabe. Class. Quant. Grav. 6, 233 (1989).

[63] N. Seiberg, E. Witten. JHEP 9909, 032 (1999).

[64] A. A. Bichl, J. M. Grimstrup, H. Grosse, L. Popp, M. Schweda, R. Wulkenhaar. JHEP 0010, 046 (2000).

[65] I. L. Buchbinder, M. Gomes, A. Yu. Petrov, V. O. Rivelles, Phys. Lett. B517, 191 (2001).

[66] A. F. Ferrari, H. O. Girotti, M. Gomes, A. Yu. Petrov, A. A. Ribeiro, V. O. Rivelles, A. J. da Silva, Phys. Rev. D69, 025008 (2004).

[67] A. F. Ferrari, H. O. Girotti, M. Gomes, A. Yu. Petrov, A. A. Ribeiro, V. O. Rivelles, A. J. da Silva, Phys. Rev. D70, 085012 (2004).

[68] A. H. Chamseddine, J. Math. Phys. 44, 2534 (2003).

[69] M. A. Cardella, D. Zanon, Class. Quant. Grav. 20, L95 (2003).

[70] N. Seiberg, 0306, 010 (2003).

[71] M. T. Grisaru, S. Penati, A. Romagnoni, Class. Quant. Grav, S1391 (2004); A. T. Banin, I. L. Buchbinder, N. G. Pletnev, JHEP 0407, 011 (2004).

[72] S. Penati, A. Romagnoni, "Covariant quantization of $N = 1/2$ SYM theory and supergauge invariance", *Preprint* hep-th/0412041.

[73] O. D. Azorkina, A. T. Banin, I. L. Buchbinder, N. G. Pletnev, "Generic chiral superfield model on anticommutative $N = 1/2$ superspace", *Preprint* hep-th/0502008.